

# Certain results on a class of Entire functions represented by Dirichlet series having complex frequencies

Niraj Kumar

Department of Mathematics, Netaji Subhas Institute of Technology, Sector 3 Dwarka, New Delhi-110078, India  
nirajkumar2001@hotmail.com

Garima Manocha

Department of Mathematics, Netaji Subhas Institute of Technology, Sector 3 Dwarka, New Delhi-110078, India  
garima89.manocha@gmail.com

---

## Abstract

Consider  $F$  to be a class of entire functions represented by Dirichlet series with complex frequencies for which  $(k!)^{c_1} e^{c_2 k |\lambda^k|} |a_k|$  is bounded. A study on certain results has been made for this set that is  $F$  is proved to be an algebra with continuous quasi-inverse, commutative Banach algebra with identity etc. Moreover, the conditions for the elements of  $F$  to possess an inverse, quasi-inverse and the form of spectrum of  $F$  are also established.

Received 11 November 2014

Accepted in final form 1 July 2015

Communicated with Vladimír Janiš.

**Keywords** Dirichlet series, Banach algebra, Quasi-inverse, Spectrum of a set.

**MSC(2010)** 30B50, 46J15, 47A10.

---

## 1 Introduction

Consider a Dirichlet series of the form

$$f(z) = \sum_{k=1}^{\infty} a_k e^{\langle \lambda^k, z \rangle}, \quad z \in \mathbb{C}^n \quad (1.1)$$

where  $\{\lambda^k\}$ ;  $\lambda^k = (\lambda_1^k, \lambda_2^k, \dots, \lambda_n^k)$ ,  $k = 1, 2, \dots$  be a sequence of complex vectors in  $\mathbb{C}^n$ . Then  $\langle \lambda^k, z \rangle = \lambda_1^k z_1 + \lambda_2^k z_2 + \dots + \lambda_n^k z_n$ . If  $a_k/s \in \mathbb{C}$  and  $\{\lambda^k\}'s$  satisfy the condition  $|\lambda^k| \rightarrow \infty$  as  $k \rightarrow \infty$  and

$$\limsup_{k \rightarrow \infty} \frac{\log |a_k|}{|\lambda^k|} = -\infty \quad (1.2)$$

$$\limsup_{k \rightarrow \infty} \frac{\log k}{|\lambda^k|} = D < \infty \quad (1.3)$$

then from [1] the Dirichlet series (1.1) represents an entire function. In this paper let  $F$  be the set of series (1.1) for which  $(k!)^{c_1} e^{c_2 k |\lambda^k|} |a_k|$  is bounded where  $c_1, c_2 \geq 0$  and

$c_1, c_2$  are simultaneously not zero. Then every element of  $F$  represents an entire function. If

$$f(z) = \sum_{k=1}^{\infty} a_k e^{\langle \lambda^k, z \rangle} \quad \text{and} \quad g(z) = \sum_{k=1}^{\infty} b_k e^{\langle \lambda^k, z \rangle}$$

define binary operations that is addition and scalar multiplication in  $F$  as

$$f(z) + g(z) = \sum_{k=1}^{\infty} (a_k + b_k) e^{\langle \lambda^k, z \rangle},$$

$$\alpha.f(z) = \sum_{k=1}^{\infty} (\alpha.a_k) e^{\langle \lambda^k, z \rangle},$$

$$f(z).g(z) = \sum_{k=1}^{\infty} \{(k!)^{c_1} e^{c_2 k |\lambda^k|} a_k b_k\} e^{\langle \lambda^k, z \rangle}.$$

The norm in  $F$  is defined as follows

$$\|f\| = \sup_{k \geq 1} (k!)^{c_1} e^{c_2 k |\lambda^k|} |a_k|. \quad (1.4)$$

If  $c_1 = c_2 = 1$  we get the norm as defined in [3] for a class of entire functions represented by Dirichlet series

$$f(s) = \sum_{n=1}^{\infty} a_n e^{\lambda_n s}, \quad s = \sigma + it, \quad (\sigma, t \in \mathbb{R}) \quad (1.5)$$

whose coefficients belonged to a commutative Banach algebra with identity and  $\lambda_n' s \in \mathbb{R}$  satisfied the condition  $0 < \lambda_1 < \lambda_2 < \lambda_3 \dots < \lambda_n \dots; \lambda_n \rightarrow \infty$  as  $n \rightarrow \infty$ . Further in the same paper authors proved the above class to be a complex FK-space and a Fréchet space. Several other results for a different class of entire Dirichlet series (1.5) may be found in [2].

In the present paper the weighted norm is generalized and various results based on the notions of Banach algebra, Quasi-inverse, Algebra with continuous quasi-inverse, Spectrum of a set have been established.

In the sequel following definitions are required to prove the main results.

**Definition 1.** A function  $g(z) \in F$  is said to be a quasi-inverse of  $f(z) \in F$  if  $f(z)*g(z) = 0$  where

$$f(z) * g(z) = f(z) + g(z) + f(z).g(z).$$

**Definition 2.** A topological algebra  $F$  is said to be an algebra with continuous quasi-inverse if there exists a neighbourhood of the zero element, every point  $f$  of which has a quasi-inverse  $f'$  and the mapping  $f \rightarrow f'$  is continuous.

**Definition 3.** The set  $\sigma(A)$  defined as

$$\sigma(A) = \{k \in K : A - kI \text{ is not invertible}\}$$

is called the spectrum of  $A$ .

## 2 Main Results

In this section main results are proved. For the definitions of terms used refer [4] and [5].

**Theorem 4.** An element  $f(z) = \sum_{k=1}^{\infty} a_k e^{\langle \lambda^k, z \rangle} \in F$  is quasi-invertible if and only if

$$\inf_{k \geq 1} \{ |1 + (k!)^{c_1} e^{c_2 k |\lambda^k|} a_k| \} > 0. \quad (2.1)$$

The quasi-inverse of  $f(z)$  is the function  $g(z) = \sum_{k=1}^{\infty} b_k e^{\langle \lambda^k, z \rangle}$  where

$$b_k = \frac{-a_k}{1 + (k!)^{c_1} e^{c_2 k |\lambda^k|} a_k}. \quad (2.2)$$

*Proof.* Let  $f(z) \in F$  be quasi-invertible. By Definition 1, there exists  $g(z) \in F$  such that  $f(z) * g(z) = 0$ . This implies

$$a_k + b_k + (k!)^{c_1} e^{c_2 k |\lambda^k|} a_k b_k = 0$$

for all  $k \geq 1$ . Let (2.1) does not hold that is

$$\inf_{k \geq 1} \{ |1 + (k!)^{c_1} e^{c_2 k |\lambda^k|} a_k| \} = 0. \quad (2.3)$$

There exists a subsequence  $\{k_t\}$  of a sequence of indices  $\{k\}$  such that  $\|f_t\| = 1$  that is

$$(k_t!)^{c_1} e^{c_2 k_t |\lambda^{k_t}|} |a_{k_t}| = 1 \text{ as } t \rightarrow \infty. \quad (2.4)$$

Thus

$$(k_t!)^{c_1} e^{c_2 k_t |\lambda^{k_t}|} |b_{k_t}| = \frac{(k_t!)^{c_1} e^{c_2 k_t |\lambda^{k_t}|} |a_{k_t}|}{|1 + (k_t!)^{c_1} e^{c_2 k_t |\lambda^{k_t}|} a_{k_t}|}$$

Using (2.3) and (2.4),

$$\|g_t\| \rightarrow \infty \text{ as } t \rightarrow \infty$$

which is a contradiction.

Conversely let (2.1) be fulfilled. The function  $g(z)$  defined by (2.2) obviously belongs to  $F$ . Thus

$$\begin{aligned} f(z) * g(z) &= \sum_{k=1}^{\infty} \{ a_k + b_k + (k!)^{c_1} e^{c_2 k |\lambda^k|} a_k b_k \} e^{\langle \lambda^k, z \rangle} \\ &= 0. \end{aligned}$$

Thus  $f(z)$  is quasi-invertible which completes the proof of the theorem.  $\square$

**Theorem 5.**  $F$  is an algebra with continuous quasi-inverse.

*Proof.* Let  $N_\epsilon(0)$  be an  $\epsilon$ -neighbourhood of 0 where  $0 < \epsilon < 1$ . Let  $p(z) \in N_\epsilon(0)$  where  $p(z) = \sum_{k=1}^{\infty} p_k e^{\langle \lambda^k, z \rangle}$ . This implies  $\|p\| < \epsilon$ . Then

$$(k!)^{c_1} e^{c_2 k |\lambda^k|} |p_k| < \epsilon \text{ for all } k \geq 1$$

which further implies

$$\inf_{k \geq 1} \{ |1 + (k!)^{c_1} e^{c_2 k |\lambda^k|} p_k| \} \geq 1 - \epsilon > 0.$$

Hence by Theorem 4,  $p(z)$  possesses a quasi-inverse say  $q(z) = \sum_{k=1}^{\infty} q_k e^{\langle \lambda^k, z \rangle}$  where

$$q_k = \frac{-p_k}{1 + (k!)^{c_1} e^{c_2 k |\lambda^k|} p_k}.$$

Now

$$\begin{aligned} \|q\| &= \sup_{k \geq 1} (k!)^{c_1} e^{c_2 k |\lambda^k|} |q_k| \\ &= \sup_{k \geq 1} \frac{(k!)^{c_1} e^{c_2 k |\lambda^k|} |p_k|}{|1 + (k!)^{c_1} e^{c_2 k |\lambda^k|} p_k|} \\ &< \frac{\epsilon}{1 - \epsilon}. \end{aligned}$$

Hence the mapping  $p(z) \rightarrow q(z)$  is continuous. Thus by Definition 2,  $F$  is an algebra with continuous quasi-inverse. Thus the theorem is proved.  $\square$

**Theorem 6.**  $F$  is a commutative Banach algebra with identity.

*Proof.* To prove the theorem we first show that  $F$  is complete under the norm defined by (1.4). Let  $\{f_{m_1}\}$  be a cauchy sequence in  $F$ . For given  $\epsilon > 0$  we find  $m$  such that

$$\|f_{m_1} - f_{m_2}\| < \epsilon \text{ where } m_1, m_2 \geq m.$$

This implies that

$$\sup_{k \geq 1} (k!)^{c_1} e^{c_2 k |\lambda^k|} |a_{m_{1k}} - a_{m_{2k}}| < \epsilon \text{ where } m_1, m_2 \geq m.$$

Clearly  $\{a_{m_{1k}}\}$  forms a cauchy sequence in the set of complex numbers for all  $k \geq 1$  and thus converges to  $a_k$ . Therefore  $f_{m_1} \rightarrow f$ . Also

$$\begin{aligned} \sup_{k \geq 1} (k!)^{c_1} e^{c_2 k |\lambda^k|} |a_k| &\leq \sup_{k \geq 1} (k!)^{c_1} e^{c_2 k |\lambda^k|} |a_{m_{1k}} - a_k| \\ &\quad + \sup_{k \geq 1} (k!)^{c_1} e^{c_2 k |\lambda^k|} |a_{m_{1k}}| \end{aligned}$$

Hence  $f(z) \in F$ . Thus  $F$  is complete. Now if  $f(z), g(z) \in F$  then

$$\begin{aligned} \|f \cdot g\| &= \sup_{k \geq 1} (k!)^{c_1} e^{c_2 k |\lambda^k|} | (k!)^{c_1} e^{c_2 k |\lambda^k|} a_k b_k | \\ &\leq \sup_{k \geq 1} (k!)^{c_1} e^{c_2 k |\lambda^k|} |a_k| \cdot \sup_{k \geq 1} (k!)^{c_1} e^{c_2 k |\lambda^k|} |b_k| \\ &= \|f\| \cdot \|g\| \end{aligned}$$

The identity element in  $F$  is

$$e(z) = \sum_{k=1}^{\infty} (k!)^{-c_1} e^{-c_2 k |\lambda^k|} e^{\langle \lambda^k, z \rangle}.$$

Hence the theorem.  $\square$

**Theorem 7.** The function  $f(z) = \sum_{k=1}^{\infty} a_k e^{\langle \lambda^k, z \rangle}$  is invertible in  $F$  if and only if

$$\{|(k!)^{-c_1} e^{-c_2 k |\lambda^k|} a_k^{-1}|\}$$

is a bounded sequence.

*Proof.* Let  $f(z)$  be invertible and  $g(z) = \sum_{k=1}^{\infty} b_k e^{\langle \lambda^k, z \rangle}$  be its inverse. Then

$$(k!)^{c_1} e^{c_2 k |\lambda^k|} a_k b_k = (k!)^{-c_1} e^{-c_2 k |\lambda^k|}$$

Equivalently

$$(k!)^{c_1} e^{c_2 k |\lambda^k|} |b_k| = |(k!)^{-c_1} e^{-c_2 k |\lambda^k|} a_k^{-1}|$$

Clearly since  $g(z) \in F$  hence

$$\{|(k!)^{-c_1} e^{-c_2 k |\lambda^k|} a_k^{-1}|\}$$

is a bounded sequence.

Conversely suppose  $\{|(k!)^{-c_1} e^{-c_2 k |\lambda^k|} a_k^{-1}|\}$  be a bounded sequence. Define  $g(z)$  such that

$$g(z) = \sum_{k=1}^{\infty} (k!)^{-2c_1} e^{-2c_2 k |\lambda^k|} a_k^{-1} e^{\langle \lambda^k, z \rangle}.$$

Obviously  $g(z) \in F$ . Moreover

$$\begin{aligned} f(z).g(z) &= \sum_{k=1}^{\infty} (k!)^{c_1} e^{c_2 k |\lambda^k|} \{a_k (k!)^{-2c_1} e^{-2c_2 k |\lambda^k|} a_k^{-1}\} e^{\langle \lambda^k, z \rangle} \\ &= e(z). \end{aligned}$$

Hence the proof of the theorem is completed. □

**Theorem 8.** The spectrum  $\sigma(f)$  where  $f(z) \in F$  is precisely of the form

$$\sigma(f) = cl\{(k!)^{c_1} e^{c_2 k |\lambda^k|} a_k : k \geq 1\}.$$

*Proof.* In Theorem 7,  $f(z) = \sum_{k=1}^{\infty} a_k e^{\langle \lambda^k, z \rangle} \in F$  is invertible if and only if

$$\{|(k!)^{-c_1} e^{-c_2 k |\lambda^k|} a_k^{-1}|\}$$

is a bounded sequence. Thus  $\{f(z) - \lambda.e(z)\}$  is not invertible if and only if

$$\{(k!)^{c_1} e^{c_2 k |\lambda^k|} |a_k - \lambda (k!)^{-c_1} e^{-c_2 k |\lambda^k|}\}^{-1}$$

is not bounded. Therefore by Definition 3, this is possible if and only if there exists a subsequence  $\{k_n\}$  of a sequence of indices  $\{k\}$  such that

$$|(k_n!)^{c_1} e^{c_2 k_n |\lambda^{k_n}|} a_{k_n} - \lambda|$$

tends to zero as  $n \rightarrow \infty$ . Equivalently

$$\lambda \in cl\{(k!)^{c_1} e^{c_2 k |\lambda^k|} a_k : k \geq 1\}$$

which proves the theorem. □

The results proved in this section would further be useful in the study of the spaces like FK-space, Fréchet space, Montel space,  $C^*$ -algebra etc. and in the study of functions preserving the asymptotic equivalence of functions and sequences that is Pseudo-regularly varying (PRV) functions. Also these results have significant applications in the fields of topology, functional analysis, modern analysis etc.

## References

- [1] L. H. Khoi, Coefficient multipliers for some classes of Dirichlet series in several complex variables, *Acta Math. Vietnamica* **24(2)** (1999), 169–182.
- [2] N. Kumar and G. Manocha, On a class of entire functions represented by Dirichlet series, *J. Egypt. Math. Soc.* **21** (2013), 21–24.
- [3] N. Kumar and G. Manocha, A class of entire Dirichlet series as an FK-space and a Frechet space, *Acta Math. Scientia* **33B(6)** (2013), 1571–1578.
- [4] R. Larsen, “Banach Algebras - An Introduction”, Marcel Dekker Inc., New York, 1973.
- [5] R. Larsen, “Functional Analysis - An Introduction”, Marcel Dekker Inc., New York, 1973.