

# $\mathcal{I}$ -derivative

Vladislav Banas

Department of Algebra, Geometry, and Mathematical Education  
Faculty of Mathematics, Physics and Informatics, Comenius University

Mlynská dolina, 842 48 Bratislava, Slovakia

[vladislav.banas@gmail.com](mailto:vladislav.banas@gmail.com)

---

## Abstract

This paper deals with a derivative of a real function based on the notion of  $\mathcal{I}$ -convergence.

Received 27 April 2013

Revised August 6, 2013

Accepted in final form September 13, 2013

Published online October 18, 2013

Communicated with Lubomír Snoha.

**Keywords**  $\mathcal{I}$ -convergence,  $\mathcal{I}^{\mathcal{K}}$ -convergence,  $\mathcal{I}$ -continuity,  $\mathcal{I}$ -derivative.

**MSC(2010)** 26A03, 26A24, 40A05.

---

## 1 Introduction

In the connection to some known results about functions preserving  $\mathcal{I}$ -convergence and  $\mathcal{I}$ -continuity (see [1, 2, 3, 4]) we introduce the  $\mathcal{I}$ -derivative of a real function, i.e. the derivative based on the notion of  $\mathcal{I}$ -convergence.

$\mathcal{I}$ -convergence was introduced in [1] as a generalization of statistical convergence (see [5, 6]).

In this paper we will elucidate the relationship of  $\mathcal{I}$ -derivative to usual derivative with respect to the choice of ideals used in the definition of  $\mathcal{I}$ -derivative.

**Definition 1.** (see [7, p. 6]) A non-void family  $\mathcal{I}$  of subsets of a given set  $X$  is called an *ideal* on  $X$  if it is hereditary and additive, i.e.

- (1)  $A \in \mathcal{I}$  and  $B \subset A \Rightarrow B \in \mathcal{I}$ ,
- (2)  $A \in \mathcal{I}$  and  $B \in \mathcal{I} \Rightarrow A \cup B \in \mathcal{I}$ .

An ideal  $\mathcal{I}$  is called a *proper* ideal if  $X \notin \mathcal{I}$ .

A proper ideal  $\mathcal{I}$  is said to be *admissible* (see [1]) if  $\mathcal{I}$  contains every singleton.

The dual notion to the notion of an ideal is the notion of a filter.

**Definition 2.** (see [7, p. 6]) A non-void family  $\mathcal{F}$  of subsets of a given set  $X$  is called a *filter* on  $X$  if

- (1)  $A \in \mathcal{F}$  and  $A \subset B \Rightarrow B \in \mathcal{F}$ ,
- (2)  $A \in \mathcal{F}$  and  $B \in \mathcal{F} \Rightarrow A \cap B \in \mathcal{F}$ .

A filter  $\mathcal{F}$  is *proper* if  $\emptyset \notin \mathcal{F}$ .

Obviously, for each ideal  $\mathcal{I}$  the system

$$\mathcal{F}(\mathcal{I}) = \{X \setminus A : A \in \mathcal{I}\}$$

is a filter on  $X$ .

**Definition 3.** (see [1]) Let  $\mathcal{I}$  be a proper ideal on the set  $\mathbb{N}$ . A sequence  $(x_n)_{n \in \mathbb{N}}$  of real numbers is said to be  $\mathcal{I}$ -convergent to  $\xi \in \mathbb{R}$  ( $\mathcal{I}\text{-lim } x_n = \xi$ ) if and only if for each  $\varepsilon > 0$  the set

$$A(\varepsilon) = \{n \in \mathbb{N} : |x_n - \xi| \geq \varepsilon\}$$

belongs to  $\mathcal{I}$ . The element  $\xi$  is called  $\mathcal{I}$ -limit of the sequence  $(x_n)_{n \in \mathbb{N}}$ .

In what follows we recall some basic properties of  $\mathcal{I}$ -convergence and of the notions  $\mathcal{I}$ -limit inferior and  $\mathcal{I}$ -limit superior (see [8, 9]).

**Theorem 4.** (see [9]) Let  $(x_n)_{n \in \mathbb{N}}$ ,  $(y_n)_{n \in \mathbb{N}}$  be sequences of real numbers such that  $\mathcal{I}\text{-lim } x_n = \xi$ ,  $\mathcal{I}\text{-lim } y_n = \eta$ . Then

- (a)  $\mathcal{I}\text{-lim}(x_n \cdot y_n) = \xi \cdot \eta$ ,
- (b)  $\mathcal{I}\text{-lim}(x_n + y_n) = \xi + \eta$ .

Let  $t \in \mathbb{R}$  and  $(x_n)_{n \in \mathbb{N}}$  be a sequence of real numbers. Put

$$M_t = \{n : x_n > t\}, \quad M^t = \{n : x_n < t\}.$$

**Definition 5.** (see [9])

- (a) If there is a  $t \in \mathbb{R}$  such that  $M_t \notin \mathcal{I}$ , we put

$$\mathcal{I}\text{-lim sup } x_n = \sup\{t \in \mathbb{R} : M_t \notin \mathcal{I}\}.$$

If  $M_t \in \mathcal{I}$  for each  $t \in \mathbb{R}$ , then  $\mathcal{I}\text{-lim sup } x_n = -\infty$ .

- (b) If there is a  $t \in \mathbb{R}$  such that  $M^t \notin \mathcal{I}$ , we put

$$\mathcal{I}\text{-lim inf } x_n = \inf\{t \in \mathbb{R} : M^t \notin \mathcal{I}\}.$$

If  $M^t \in \mathcal{I}$  for each  $t \in \mathbb{R}$ , then  $\mathcal{I}\text{-lim inf } x_n = +\infty$ .

**Theorem 6.** (see [9]) The inequality

$$\mathcal{I}\text{-lim inf } x_n \leq \mathcal{I}\text{-lim sup } x_n$$

holds for every sequence  $(x_n)_{n \in \mathbb{N}}$  of real numbers.

**Theorem 7.** (see [9]) The sequence  $(x_n)_{n \in \mathbb{N}}$  of real numbers is  $\mathcal{I}$ -convergent if and only if

$$\mathcal{I}\text{-lim inf } x_n = \mathcal{I}\text{-lim sup } x_n.$$

If this equality holds, then

$$\mathcal{I}\text{-lim } x_n = \mathcal{I}\text{-lim inf } x_n = \mathcal{I}\text{-lim sup } x_n.$$

We use various types of convergences in the article. We also often use sequences of real numbers with restrictions on their members. In order to keep the text comprehensible we introduce following notations.

Let  $\mathbb{R}^{\mathbb{N}}$  denote the set of all sequences of real numbers. Let  $x = (x_n)_{n \in \mathbb{N}}$ . We put

$$\mathcal{S} = \{x \in \mathbb{R}^{\mathbb{N}} : x_n \neq 0, n \in \mathbb{N}\},$$

$$\mathcal{S}_{\mathcal{I}} = \{x : \mathcal{I}\text{-}\lim x_n = 0, x_n \neq 0, n \in \mathbb{N}\}$$

and

$$\mathcal{S}_{\mathcal{I}}^+ = \{x \in \mathcal{S}_{\mathcal{I}} : x_n > 0, n \in \mathbb{N}\}, \mathcal{S}_{\mathcal{I}}^- = \{x \in \mathcal{S}_{\mathcal{I}} : x_n < 0, n \in \mathbb{N}\}.$$

Note that  $\mathcal{S}_{\mathcal{I}_f}$ , where  $\mathcal{I}_f$  is the Fréchet ideal, contains only sequences convergent in the usual sense.

## 2 $\mathcal{I}$ -derivative

**Definition 8.** Let  $\mathcal{I}$  be an admissible ideal on the set  $\mathbb{N}$ . A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  has an  $\mathcal{I}$ -derivative  $d \in \mathbb{R}$  at a point  $x_0$ , i.e.  $\mathcal{I}\text{-}f'(x_0) = d$ , if and only if

$$\mathcal{I}\text{-}\lim \frac{f(x_0 + x_n) - f(x_0)}{x_n} = d \quad (2.1)$$

holds for each sequence  $(x_n)_{n \in \mathbb{N}} \in \mathcal{S}_{\mathcal{I}}$ .

In [2, Theorem 1] it is showed that if a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is  $\mathcal{I}$ -continuous at a point  $x_0$ , i.e.

$$\mathcal{I}\text{-}\lim x_n = x_0 \Rightarrow \mathcal{I}\text{-}\lim f(x_n) = f(x_0) \quad (2.2)$$

holds for each sequence  $(x_n)_{n \in \mathbb{N}}$ , then it is continuous at the point  $x_0$ .

**Proposition 9.** Let a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  have an  $\mathcal{I}$ -derivative  $d \in \mathbb{R}$  at a point  $x_0$  where  $\mathcal{I}$  is an admissible ideal on the set  $\mathbb{N}$ . Then  $f$  is continuous at the point  $x_0$ .

*Proof.* Let  $(x_n)_{n \in \mathbb{N}} \in \mathcal{S}_{\mathcal{I}}$ . Obviously

$$f(x_0 + x_n) - f(x_0) = \frac{f(x_0 + x_n) - f(x_0)}{x_n} \cdot x_n$$

holds for each  $n \in \mathbb{N}$ . Thus

$$\mathcal{I}\text{-}\lim(f(x_0 + x_n) - f(x_0)) = \mathcal{I}\text{-}\lim \frac{f(x_0 + x_n) - f(x_0)}{x_n} \cdot x_n.$$

According to the assumption and Theorem 4 we have

$$\mathcal{I}\text{-}\lim f(x_0 + x_n) = f(x_0)$$

what means that (2.2) holds for each sequence of real numbers and so  $f$  is continuous at the point  $x_0$ .  $\square$

For  $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $x_0 \in \mathbb{R}$  put

$$D^+ f(x_0) = \limsup_{h \rightarrow 0^+} \frac{f(x_0 + h) - f(x_0)}{h},$$

$$D_+ f(x_0) = \liminf_{h \rightarrow 0^+} \frac{f(x_0 + h) - f(x_0)}{h},$$

$$D^- f(x_0) = \limsup_{h \rightarrow 0^-} \frac{f(x_0 + h) - f(x_0)}{h},$$

$$D_- f(x_0) = \liminf_{h \rightarrow 0^-} \frac{f(x_0 + h) - f(x_0)}{h}.$$

The numbers  $D^+ f(x_0), D_+ f(x_0), D^- f(x_0), D_- f(x_0)$  are called Dini's derivatives (see e.g. [10, p. 27]) and it is well known that all Dini's derivatives at the point  $x_0$  are equal to  $d \in \mathbb{R}$  if and only if  $f'(x_0) = d$ .

For each  $x \in \mathcal{S}_{\mathcal{I}}^+$  we put

$$D_x^+ f(x_0) = \mathcal{I}\text{-}\limsup \frac{f(x_0 + x_n) - f(x_0)}{x_n},$$

$$D_+^x f(x_0) = \mathcal{I}\text{-}\liminf \frac{f(x_0 + x_n) - f(x_0)}{x_n},$$

and for each  $y \in \mathcal{S}_{\mathcal{I}}^-$

$$D_y^- f(x_0) = \mathcal{I}\text{-}\limsup \frac{f(x_0 + y_n) - f(x_0)}{y_n},$$

$$D_y^y f(x_0) = \mathcal{I}\text{-}\liminf \frac{f(x_0 + y_n) - f(x_0)}{y_n}.$$

**Lemma 10.** *Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $x_0 \in \mathbb{R}$ . Let  $(x_n)_{n \in \mathbb{N}}$  be an arbitrary sequence of real numbers such that  $\mathcal{I}\text{-}\lim x_n = x_0$ ,  $x_n \neq x_0$  for each  $n \in \mathbb{N}$  where  $\mathcal{I}$  is an admissible ideal on the set  $\mathbb{N}$ . Then*

$$\liminf_{x \rightarrow x_0} f(x) \leq \mathcal{I}\text{-}\liminf f(x_n), \quad (2.3)$$

$$\mathcal{I}\text{-}\limsup f(x_n) \leq \limsup_{x \rightarrow x_0} f(x). \quad (2.4)$$

*Proof.* We will prove (2.3), the proof of (2.4) is quite similar.

Let  $\lambda > 0$ . Put  $O_\lambda = \{f(x) : 0 < |x - x_0| < \lambda\}$ . Because  $(x_n)_{n \in \mathbb{N}}$  is  $\mathcal{I}$ -convergent, the set  $\{n : |x_n - x_0| \geq \lambda\}$  belongs to  $\mathcal{I}$  what implies  $\{n : f(x_n) \in \mathbb{R} \setminus O_\lambda\} \in \mathcal{I}$ . Let  $s = \inf O_\lambda$ . Case  $s = -\infty$  is trivial so suppose that  $s \in \mathbb{R}$ . It is obvious that  $\{f(x_n) : f(x_n) < s\} \subset \mathbb{R} \setminus O_\lambda$  and so  $\{n : f(x_n) < s\}$  belongs to  $\mathcal{I}$  what implies  $s \notin \{t \in \mathbb{R} : \{n : f(x_n) < t\} \notin \mathcal{I}\}$ . From the arbitrariness of the choice of  $\lambda$  we have

$$\sup_{\lambda > 0} \{\inf O_\lambda\} \leq \inf \{t \in \mathbb{R} : \{n : f(x_n) < t\} \notin \mathcal{I}\}$$

and the statement holds.  $\square$

**Theorem 11.** *Let  $\mathcal{I}$  be an admissible ideal on the set  $\mathbb{N}$ . A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  has a derivative  $d \in \mathbb{R}$  at a point  $x_0$  if and only if  $\mathcal{I}\text{-}f'(x_0) = d$ .*

*Proof.* Let  $f'(x_0) = d$ . According to Theorem 6, Theorem 7 and Lemma 10 for each  $x \in \mathcal{S}_{\mathcal{I}}^+$  we have

$$D_+ f(x_0) \leq D_+^x f(x_0) \leq D_x^+ f(x_0) \leq D^+ f(x_0)$$

and for each  $y \in \mathcal{S}_{\mathcal{I}}^-$

$$D_- f(x_0) \leq D_y^y f(x_0) \leq D_y^- f(x_0) \leq D^- f(x_0).$$

Therefore (2.1) holds for each sequence from  $\mathcal{S}_{\mathcal{I}}^+ \cup \mathcal{S}_{\mathcal{I}}^-$ . It is sufficient to show, that (2.1) holds for each  $z \in \mathcal{S}_{\mathcal{I}} \setminus (\mathcal{S}_{\mathcal{I}}^+ \cup \mathcal{S}_{\mathcal{I}}^-)$ . By contradiction. Let there exists  $z \in \mathcal{S}_{\mathcal{I}} \setminus (\mathcal{S}_{\mathcal{I}}^+ \cup \mathcal{S}_{\mathcal{I}}^-)$  such that (2.1) does not hold. For  $\eta > 0$  we put

$$K_{\eta} = \left\{ n : \frac{f(x_0 + z_n) - f(x_0)}{z_n} \notin B(d, \eta) \right\},$$

where  $B(d, \eta)$  is open ball with center  $d$  and radius  $\eta$ . There is  $\eta_0$  such that  $K_{\eta_0} \notin \mathcal{I}$ . Hence at least one of sets

$$A = \{n : n \in K_{\eta_0}, z_n < 0\}, B = \{n : n \in K_{\eta_0}, z_n > 0\}$$

does not belong to  $\mathcal{I}$  and it is infinite. Let it be the set  $B$ . Define the sequence  $s = (s_n)_{n \in \mathbb{N}}$  as follows. For  $n \in B$  put  $s_n = z_n$  and  $s_n = \frac{1}{n}$  if  $n \in \mathbb{N} \setminus B$ . Obviously  $s \in \mathcal{S}_{\mathcal{I}}^+ \cup \mathcal{S}_{\mathcal{I}}^-$ , a contradiction. If the set  $A$  does not belong to  $\mathcal{I}$  for  $n \in A$  we put  $s_n = z_n$  and  $s_n = -\frac{1}{n}$  if  $n \in \mathbb{N} \setminus A$ .

Suppose now, that  $f$  does not have the derivative at the point  $x_0$ . This implication follows immediately from the fact, that if one of Dini's derivatives is equal to  $s \in \mathbb{R}$ , there is a sequence  $(x_n)_{n \in \mathbb{N}}$  convergent to  $x_0$  such that the sequence  $(y_n)_{n \in \mathbb{N}}$  of numbers

$$y_n = \frac{f(x_0 + x_n) - f(x_0)}{x_n}$$

converges to  $s$  or if  $s = +\infty(-\infty)$  there is a sequence  $(x_n)_{n \in \mathbb{N}}$  convergent to  $x_0$  such that the sequence  $(y_n)_{n \in \mathbb{N}}$  is increasing (decreasing) with limit  $+\infty(-\infty)$ . Hence for each admissible ideal  $\mathcal{I}$  the function  $f$  does not have  $\mathcal{I}$ -derivative at the point  $x_0$ .  $\square$

In Definition 8 we have used the same ideal  $\mathcal{I}$  for  $\mathcal{I}$ -convergence in  $\mathcal{S}_{\mathcal{I}}$  and in (2.1). It is quite natural to ask what (if any) difference in our results will be reached by using various ideals.

**Definition 12.** Let  $\mathcal{I}_1, \mathcal{I}_2$  be admissible ideals on the set  $\mathbb{N}$ . A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  has a  $(\mathcal{I}_1, \mathcal{I}_2)$ -derivative  $d \in \mathbb{R}$  at a point  $x_0$ , i.e.  $(\mathcal{I}_1, \mathcal{I}_2)$ - $f'(x_0) = d$ , if and only if

$$\mathcal{I}_1\text{-}\lim x_n = 0 \Rightarrow \mathcal{I}_2\text{-}\lim \frac{f(x_0 + x_n) - f(x_0)}{x_n} = d \quad (2.5)$$

holds for each sequence  $(x_n)_{n \in \mathbb{N}} \in \mathcal{S}$ .

**Remark 13.** Studying the proof of Theorem 11 we find that in case  $\mathcal{I}_1 \subset \mathcal{I}_2$  we get the same results for  $(\mathcal{I}_1, \mathcal{I}_2)$ -derivative as well. The difference is reached if  $\mathcal{I}_1 \setminus \mathcal{I}_2 \neq \emptyset$ . The following theorem says that in this case  $(\mathcal{I}_1, \mathcal{I}_2)$ -derivative is no longer only local property of a real function at a point.

**Theorem 14.** Let  $\mathcal{I}_1, \mathcal{I}_2$  be admissible ideals on the set  $\mathbb{N}$  such that  $\mathcal{I}_1 \setminus \mathcal{I}_2 \neq \emptyset$ . Then a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  has a  $(\mathcal{I}_1, \mathcal{I}_2)$ -derivative  $d \in \mathbb{R}$  at a point  $x_0$  if and only if  $f$  is a linear function.

*Proof.* The case if  $f$  is a linear function is trivial.

Suppose that  $f$  is not linear and  $(\mathcal{I}_1, \mathcal{I}_2)$ - $f'(x_0) = d$ . So there exists  $z \in \mathbb{R} \setminus \{0\}$  such that

$$\frac{f(x_0 + z) - f(x_0)}{z} = d' \neq d. \quad (2.6)$$

Because  $\mathcal{I}_1, \mathcal{I}_2$  are both admissible, there is an infinite set  $A \in \mathcal{I}_1 \setminus \mathcal{I}_2$ . Define sequence  $(x_n)_{n \in \mathbb{N}}$  as follows. If  $n \in A$  put  $x_n = z$  else  $x_n = \frac{1}{n}$ . Obviously  $\mathcal{I}_1\text{-lim } x_n = 0$ . Let  $\eta > 0$  be such that  $d' \notin B(d, \eta)$ . Then

$$\left\{ n : \frac{f(x_0 + x_n) - f(x_0)}{x_n} \notin B(d, \eta) \right\} \notin \mathcal{I}_2.$$

That is a contradiction because for the sequence  $(x_n)_{n \in \mathbb{N}}$  the implication (2.5) does not hold.  $\square$

Simultaneously with  $\mathcal{I}$ -convergence another closely related kind of convergence called  $\mathcal{I}^*$ -convergence was introduced and investigated in [1, 9, 3], later generalized as  $\mathcal{I}^{\mathcal{K}}$ -convergence in [11].

Let  $\mathcal{I}$  be an ideal on a set  $S$  and  $X$  be a topological space. A function  $f : S \rightarrow X$  is said to be  $\mathcal{I}$ -convergent to  $x \in X$  if

$$f^{-1}(U) = \{s \in S : f(s) \in U\} \in \mathcal{F}(\mathcal{I})$$

holds for each neighborhood of  $x$ .

**Definition 15.** (see [11]) Let  $\mathcal{I}, \mathcal{K}$  be ideals on a set  $S$ . Let  $X$  be a topological space and  $x \in X$ . A function  $f : S \rightarrow X$  is said to be  $\mathcal{I}^{\mathcal{K}}$ -convergent to  $x$  if there exists a set  $M \in \mathcal{F}(\mathcal{I})$  such that the function  $g : S \rightarrow X$  given by

$$g(s) = \begin{cases} f(s), & \text{if } s \in M \\ x, & \text{otherwise} \end{cases}$$

is  $\mathcal{K}$ -convergent to  $x$ .

**Proposition 16.** Let  $\mathcal{I}$  be an arbitrary and  $\mathcal{K}$  an admissible ideal on the set  $\mathbb{N}$ . Let  $(y_n)_{n \in \mathbb{N}} \in \mathbb{R}^{\mathbb{N}}$ ,  $\lim y_n = x$ . Then a real sequence  $(x_n)_{n \in \mathbb{N}}$  is  $\mathcal{I}^{\mathcal{K}}$ -convergent to  $x$  if and only if there is a set  $M \in \mathcal{F}(\mathcal{I})$  such that the sequence  $(z_n)_{n \in \mathbb{N}}$ ,  $z_n = x_n$  for  $n \in M$  otherwise  $z_n = y_n$ , is  $\mathcal{K}$ -convergent to  $x$ .

*Proof.* Let  $(x_n)_{n \in \mathbb{N}}$  be  $\mathcal{I}^{\mathcal{K}}$ -convergent to  $x$ ,  $\eta > 0$ . Let  $M$  be the corresponding set belonging to  $\mathcal{F}(\mathcal{I})$ . Then  $U = \{n : z_n \notin B(x, \eta) \wedge n \in M\}$  belongs to  $\mathcal{K}$ . The set  $V = \{n : z_n \notin B(x, \eta) \wedge n \notin M\}$  is always finite or empty and since  $\mathcal{K}$  is admissible the union  $U \cup V$  belongs to  $\mathcal{K}$  as well.

The converse implication can be proved using similar consideration.  $\square$

In [11] it is showed that  $\mathcal{I}^{\mathcal{K}}$ -convergence implies  $\mathcal{I}$ -convergence if  $\mathcal{K} \subset \mathcal{I}$ . The converse implication holds (assuming that  $X$  is a first countable topological space) if  $\mathcal{I}$  has additive property with respect to  $\mathcal{K}$ , or more briefly that the condition  $AP(\mathcal{I}, \mathcal{K})$  holds. The condition  $AP(\mathcal{I}, \mathcal{K})$  holds, if for every sequence of mutually disjoint sets  $(A_n)_{n \in \mathbb{N}}$  belonging to  $\mathcal{I}$  there is a sequence  $(B_n)_{n \in \mathbb{N}}$  of sets belonging to  $\mathcal{I}$  such that  $A_n \triangle B_n \in \mathcal{K}$  for  $n \in \mathbb{N}$  and  $B = \bigcup_{n \in \mathbb{N}} B_n \in \mathcal{I}$ .

**Definition 17.** Let  $\mathcal{I}, \mathcal{K}$  be admissible ideals on the set  $\mathbb{N}$ .

A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  has an  $\mathcal{I}^{\mathcal{K}}$ -derivative  $d \in \mathbb{R}$  at a point  $x_0$ , i.e.  $\mathcal{I}^{\mathcal{K}}\text{-}f'(x_0) = d$ , if and only if

$$\mathcal{I}^{\mathcal{K}}\text{-lim } x_n = 0 \Rightarrow \mathcal{I}^{\mathcal{K}}\text{-lim } \frac{f(x_0 + x_n) - f(x_n)}{x_n} = d$$

holds for each sequence  $(x_n)_{n \in \mathbb{N}} \in \mathcal{S}$ .

**Theorem 18.** Let  $\mathcal{I}, \mathcal{K}$  be admissible ideals on the set  $\mathbb{N}$  and  $f : \mathbb{R} \rightarrow \mathbb{R}$ .

(a) If  $f'(x_0) = d$ , then  $\mathcal{I}^{\mathcal{K}}-f'(x_0) = d$ .

(b) If  $\mathcal{I}$  fulfils condition  $AP(\mathcal{I}, \mathcal{K})$ ,  $\mathcal{K} \subset \mathcal{I}$  and  $\mathcal{I}^{\mathcal{K}}-f'(x_0) = d$ , then  $f'(x_0) = d$ .

*Proof.* (a) Let  $(x_n)_{n \in \mathbb{N}} \in \mathcal{S}_{\mathcal{I}^{\mathcal{K}}}$ ,  $(y_n)_{n \in \mathbb{N}} \in \mathcal{S}_{\mathcal{I}_f}$ . Thus there is a set  $M \in \mathcal{F}(\mathcal{I})$  such that the sequence  $(z_n)_{n \in \mathbb{N}}$ ,  $z_n = x_n$  if  $n \in M$  else  $z_n = y_n$ , is  $\mathcal{K}$ -convergent to the point 0. According to assumption, Proposition 16 and Theorem 11 we have

$$\mathcal{I}^{\mathcal{K}}\text{-}\lim \frac{f(x_0 + x_n) - f(x_n)}{x_n} = d.$$

(b) Let  $(x_n)_{n \in \mathbb{N}} \in \mathcal{S}_{\mathcal{I}}$ . Because  $\mathcal{I}$  fulfils condition  $AP(\mathcal{I}, \mathcal{K})$  and  $\mathcal{K} \subset \mathcal{I}$  the implications

$$\begin{aligned} \mathcal{I}\text{-}\lim x_n = 0 &\Rightarrow \mathcal{I}^{\mathcal{K}}\text{-}\lim x_n = 0, \\ \mathcal{I}^{\mathcal{K}}\text{-}\lim \frac{f(x_0 + x_n) - f(x_n)}{x_n} = d &\Rightarrow \mathcal{I}\text{-}\lim \frac{f(x_0 + x_n) - f(x_n)}{x_n} = d \end{aligned}$$

hold. Therefore  $\mathcal{I}\text{-}f'(x_0) = d$  what is equivalent (Theorem 11) with  $f'(x_0) = d$ .  $\square$

According to Definition 12 we can introduce next definition.

**Definition 19.** Let  $\mathcal{I}_1, \mathcal{I}_2, \mathcal{K}_1, \mathcal{K}_2$  be admissible ideals on the set  $\mathbb{N}$ .

A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  has a  $(\mathcal{I}_1^{\mathcal{K}_1}, \mathcal{I}_2^{\mathcal{K}_2})$ -derivative  $d \in \mathbb{R}$  at a point  $x_0$ , i.e.  $(\mathcal{I}_1^{\mathcal{K}_1}, \mathcal{I}_2^{\mathcal{K}_2})\text{-}f'(x_0) = d$ , if and only if

$$\mathcal{I}_1^{\mathcal{K}_1}\text{-}\lim x_n = 0 \Rightarrow \mathcal{I}_2^{\mathcal{K}_2}\text{-}\lim \frac{f(x_0 + x_n) - f(x_n)}{x_n} = d$$

holds for each sequence  $(x_n)_{n \in \mathbb{N}} \in \mathcal{S}$ .

**Theorem 20.** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $\mathcal{I}_1, \mathcal{I}_2, \mathcal{K}_1, \mathcal{K}_2$  be admissible ideals on the set  $\mathbb{N}$ . Let  $\mathcal{I}_1, \mathcal{I}_2$  possess conditions  $AP(\mathcal{I}_1, \mathcal{K}_1)$ ,  $AP(\mathcal{I}_2, \mathcal{K}_2)$  and  $\mathcal{K}_1 \subset \mathcal{I}_1$ ,  $\mathcal{K}_2 \subset \mathcal{I}_2$ .

(a) If  $\mathcal{I}_1 \subset \mathcal{I}_2$  then  $(\mathcal{I}_1^{\mathcal{K}_1}, \mathcal{I}_2^{\mathcal{K}_2})\text{-}f'(x_0) = d$  if and only if  $f'(x_0) = d$ .

(b) If  $\mathcal{I}_1 \setminus \mathcal{I}_2 \neq \emptyset$  then  $(\mathcal{I}_1^{\mathcal{K}_1}, \mathcal{I}_2^{\mathcal{K}_2})\text{-}f'(x_0) = d$  if and only if  $f$  is a linear function.

*Proof.* According to assumption the following equivalences

$$\begin{aligned} \mathcal{I}_1\text{-}\lim x_n = 0 &\Leftrightarrow \mathcal{I}_1^{\mathcal{K}_1}\text{-}\lim x_n = 0, \\ \mathcal{I}_2\text{-}\lim \frac{f(x_0 + x_n) - f(x_n)}{x_n} = d &\Leftrightarrow \mathcal{I}_2^{\mathcal{K}_2}\text{-}\lim \frac{f(x_0 + x_n) - f(x_n)}{x_n} = d \end{aligned}$$

hold. The statement of Theorem 20 follows from Remark 13 and Theorem 14.  $\square$

## References

- [1] P. Kostyrko, T. Šalát and W. Wilczyński,  $\mathcal{I}$ -convergence, *Real Anal. Exchange* **26(2)** (2000-2001), 669–686.
- [2] V. Baláž, J. Červeňanský, P. Kostyrko and T. Šalát,  $\mathcal{I}$ -convergence and  $\mathcal{I}$ -continuity of real functions, *Acta Mathematica, Faculty of Natural Sciences, Constantine the Philosopher University Nitra* **5** (2002), 43–50.
- [3] B. K. Lahiri and P. Das,  $\mathcal{I}$  and  $\mathcal{I}^*$  convergence in topological spaces, *Math. Bohem.* **2** (2005), 153–160.

- [4] M. Sleziak,  $\mathcal{I}$ -continuity in topological spaces, *Acta Mathematica, Faculty of Natural Sciences, Constantine the Philosopher University Nitra* **6** (2003), 115–122.
- [5] H. Fast, Sur la convergence statistique, *Colloq. Math.* **2** (1951), 241–244.
- [6] I. J. Schoenberg, The integrability of certain functions and related summability methods, *Amer. Math. Monthly* **66** (1959), 361–375.
- [7] K. Kuratowski, “Topologie, Vol. I.”, New edition, revised and augmented, PWN, Warszawa, 1966.
- [8] K. Demirci,  $\mathcal{I}$ -limit superior and limit inferior, *Math. Commun.* **6** (2001), 165–172.
- [9] P. Kostyrko, M. Mačaj, M. Sleziak and T. Šalát,  $\mathcal{I}$ -convergence and extremal  $\mathcal{I}$ -limit points, *Math. Slovaca* **55(4)** (2005), 443–464.
- [10] A. C. M. van Rooij and W. H. Schikhof, “A Second Course on Real Functions”, Cambridge University Press, Cambridge, 1982.
- [11] M. Mačaj and M. Sleziak,  $\mathcal{I}^{\mathcal{K}}$ -convergence, *Real Anal. Exchange* **36(1)** (2010-2011), 177–194.