

Some Perturbed Ostrowski Type Inequalities for Absolutely Continuous Functions (I)

Sever S. Dragomir

College of Engineering & Science
Victoria University, P.O. Box 14428
Melbourne City, MC 8001, Australia.
Sever.Dragomir@vu.edu.au

Abstract

In this paper, some two parameters perturbed Ostrowski type inequalities for absolutely continuous functions are established.

Received 23 February 2015

Accepted in final form 25 March 2015

Communicated with Lubomír Snoha.

Keywords Ostrowski's inequality, Integral inequalities, Perturbed Ostrowski Type Inequalities..

MSC(2010) 25D15, 25D10.

1 Introduction

We start with the following result that generalizes Ostrowski's inequality for real valued differentiable functions whose derivative are bounded.

Theorem 1 (Dragomir, 2003 [20]). *Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$ and $x \in [a, b]$. Suppose that there exist the functions $m_i, M_i : [a, b] \rightarrow \mathbb{R}$ ($i = \overline{1, 2}$) with the properties:*

$$m_1(x) \leq f'(t) \leq M_1(x) \quad \text{for a.e. } t \in [a, x] \quad (1.1)$$

and

$$m_2(x) \leq f'(t) \leq M_2(x) \quad \text{for a.e. } t \in (x, b]. \quad (1.2)$$

Then we have the inequalities:

$$\begin{aligned} & \frac{1}{2(b-a)} \left[m_1(x)(x-a)^2 - M_2(x)(b-x)^2 \right] \\ & \leq f(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ & \leq \frac{1}{2(b-a)} \left[M_1(x)(x-a)^2 - m_2(x)(b-x)^2 \right]. \end{aligned} \quad (1.3)$$

The constant $\frac{1}{2}$ is sharp on both sides.

In the case that the derivative is globally bounded on $[a, b]$ by two constants, then we have:

Corollary 2. *If $f : [a, b] \rightarrow \mathbb{R}$ is absolutely continuous on $[a, b]$ and the derivative $f' : [a, b] \rightarrow \mathbb{R}$ is bounded above and below, that is, there exists the constants $M > m$ such that*

$$-\infty < m \leq f'(t) \leq M < \infty \text{ for a.e. } t \in [a, b], \quad (1.4)$$

then we have the inequality

$$\begin{aligned} & \frac{1}{2(b-a)} \left[m(x-a)^2 - M(b-x)^2 \right] \\ & \leq f(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ & \leq \frac{1}{2(b-a)} \left[M(x-a)^2 - m(b-x)^2 \right] \end{aligned} \quad (1.5)$$

for all $x \in [a, b]$. The constant $\frac{1}{2}$ is the best in both inequalities.

We may rewrite Corollary 2 in the following equivalent manner:

Corollary 3. *With the assumptions on Corollary 2, we have:*

$$\begin{aligned} & \left| f(x) - \left(x - \frac{a+b}{2} \right) \left(\frac{M+m}{2} \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{2} (M-m)(b-a) \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] \end{aligned} \quad (1.6)$$

for all $x \in [a, b]$.

Remark 4. If we assume that $\|f'\|_\infty := \operatorname{ess\,sup}_{t \in [a, b]} |f'(t)| < \infty$, then obviously we may choose in (1.5) $m = \|f'\|_\infty$ and $M = \|f'\|_\infty$, obtaining Ostrowski's inequality for absolutely continuous functions whose derivatives are essentially bounded:

$$\begin{aligned} \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| & \leq \frac{\|f'\|_\infty}{2(b-a)} \left[(x-a)^2 + (b-x)^2 \right] \\ & = \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] (b-a) \|f'\|_\infty, \end{aligned}$$

for all $x \in [a, b]$. The constant $\frac{1}{4}$ here is best.

Remark 5. Ostrowski's inequality for absolutely continuous mappings in terms of $\|f'\|_\infty$ basically states that

$$\begin{aligned} -\frac{\|f'\|_\infty}{2(b-a)} \left[(x-a)^2 + (b-x)^2 \right] & \leq f(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ & \leq \frac{\|f'\|_\infty}{2(b-a)} \left[(x-a)^2 + (b-x)^2 \right] \end{aligned} \quad (1.7)$$

for all $x \in [a, b]$.

Now, if we assume that (1.1) and (1.2) hold, then $-\|f'\|_\infty \leq m_1(x)$, $m_2(x)$ and $M_1(x)$, $M_2(x) \leq \|f'\|_\infty$, which implies:

$$\begin{aligned} & -\frac{\|f'\|_\infty}{2(b-a)} \left[(x-a)^2 + (b-x)^2 \right] \\ & \leq \frac{1}{2(b-a)} \left[m_1(x)(x-a)^2 - M_2(x)(b-x)^2 \right] \leq f(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ & \leq \frac{1}{2(b-a)} \left[M_1(x)(x-a)^2 - m_2(x)(b-x)^2 \right] \\ & \leq \frac{\|f'\|_\infty}{2(b-a)} \left[(x-a)^2 + (b-x)^2 \right]. \end{aligned} \quad (1.8)$$

Thus, the inequality (1.3) may also be regarded as a refinement of the classical Ostrowski result.

An important particular case is $x = \frac{a+b}{2}$ providing the following corollary.

Corollary 6. Assume that the derivative $f' : [a, b] \rightarrow \mathbb{R}$ satisfy the conditions:

$$-\infty < m_1 \leq f'(t) \leq M_1 < \infty \text{ for a.e. } t \in \left[a, \frac{a+b}{2} \right] \quad (1.9)$$

and

$$-\infty < m_2 \leq f'(t) \leq M_2 < \infty \text{ for a.e. } t \in \left(\frac{a+b}{2}, b \right]. \quad (1.10)$$

Then we have the inequalities

$$\begin{aligned} \frac{1}{8}(m_1 - M_2)(b-a) & \leq f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \\ & \leq \frac{1}{8}(M_1 - m_2)(b-a). \end{aligned} \quad (1.11)$$

The constant $\frac{1}{8}$ is the best in both inequalities.

Finally, if we know some global bounds for the derivative f' on $[a, b]$, then we may state the following corollary.

Corollary 7. Under the assumptions of Corollary 2, we have the midpoint inequality:

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{8}(M-m)(b-a). \quad (1.12)$$

The constant $\frac{1}{8}$ is best.

For other Ostrowski type inequalities see [1]-[19] and [21]-[42].

Motivated by the above results, we establish in this paper some perturbed Ostrowski type inequalities for complex valued differentiable functions whose derivatives are either bounded or of bounded variation. Applications for midpoint inequalities are provided as well.

2 Some Identities

We start with the following identity that will play an important role in the following:

Lemma 8. *Let $f : [a, b] \rightarrow \mathbb{C}$ be an absolutely continuous on $[a, b]$ and $x \in [a, b]$. Then for any $\lambda_1(x)$ and $\lambda_2(x)$ complex numbers, we have*

$$\begin{aligned} f(x) + \frac{1}{2(b-a)} \left[(b-x)^2 \lambda_2(x) - (x-a)^2 \lambda_1(x) \right] - \frac{1}{b-a} \int_a^b f(t) dt \\ = \frac{1}{b-a} \int_a^x (t-a) [f'(t) - \lambda_1(x)] dt + \frac{1}{b-a} \int_x^b (t-b) [f'(t) - \lambda_2(x)] dt, \end{aligned} \quad (2.1)$$

where the integrals in the right hand side are taken in the Lebesgue sense.

Proof. Utilising the integration by parts formula in the Lebesgue integral, we have

$$\begin{aligned} \int_a^x (t-a) [f'(t) - \lambda_1(x)] dt \\ = (t-a) [f(t) - \lambda_1(x)t] \Big|_a^x - \int_a^x [f(t) - \lambda_1(x)t] dt \\ = (x-a) [f(x) - \lambda_1(x)x] - \int_a^x f(t) dt + \frac{1}{2} \lambda_1(x) (x^2 - a^2) \\ = (x-a) f(x) - \lambda_1(x)x(x-a) - \int_a^x f(t) dt + \frac{1}{2} \lambda_1(x) (x^2 - a^2) \\ = (x-a) f(x) - \int_a^x f(t) dt - \frac{1}{2} (x-a)^2 \lambda_1(x) \end{aligned} \quad (2.2)$$

and

$$\begin{aligned} \int_x^b (t-b) [f'(t) - \lambda_2(x)] dt \\ = (t-b) [f(t) - \lambda_2(x)t] \Big|_x^b - \int_x^b [f(t) - \lambda_2(x)t] dt \\ = (b-x) [f(x) - \lambda_2(x)x] - \int_x^b f(t) dt + \frac{1}{2} \lambda_2(x) (b^2 - x^2) \\ = (b-x) f(x) - \int_x^b f(t) dt - (b-x) \lambda_2(x)x + \frac{1}{2} \lambda_2(x) (b^2 - x^2) \\ = (b-x) f(x) - \int_x^b f(t) dt + \frac{1}{2} (b-x)^2 \lambda_2(x). \end{aligned} \quad (2.3)$$

If we add the identities (2.2) and (2.3) and divide by $b-a$ we deduce the desired identity (2.1). \square

Corollary 9. *With the assumption in Lemma 8, we have for any $\lambda(x) \in \mathbb{C}$ that*

$$\begin{aligned} f(x) + \left(\frac{a+b}{2} - x \right) \lambda(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ = \frac{1}{b-a} \int_a^x (t-a) [f'(t) - \lambda(x)] dt + \frac{1}{b-a} \int_x^b (t-b) [f'(t) - \lambda(x)] dt. \end{aligned} \quad (2.4)$$

Remark 10. If we take $\lambda(x) = 0$ in (2.4), then we get Montgomery's identity for absolutely continuous functions, i.e.

$$\begin{aligned} f(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ = \frac{1}{b-a} \int_a^x (t-a) f'(t) dt + \frac{1}{b-a} \int_x^b (t-b) f'(t) dt, \end{aligned} \quad (2.5)$$

for $x \in [a, b]$.

We have the following midpoint representation:

Corollary 11. *With the assumption in Lemma 8, we have for any $\lambda_1, \lambda_2 \in \mathbb{C}$ that*

$$\begin{aligned} f\left(\frac{a+b}{2}\right) + \frac{1}{8}(b-a)(\lambda_2 - \lambda_1) - \frac{1}{b-a} \int_a^b f(t) dt \\ = \frac{1}{b-a} \int_a^{\frac{a+b}{2}} (t-a) [f'(t) - \lambda_1] dt + \frac{1}{b-a} \int_{\frac{a+b}{2}}^b (t-b) [f'(t) - \lambda_2] dt. \end{aligned} \quad (2.6)$$

In particular, if $\lambda_1 = \lambda_2 = \lambda$, then we have the equality

$$\begin{aligned} f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \\ = \frac{1}{b-a} \int_a^{\frac{a+b}{2}} (t-a) [f'(t) - \lambda] dt + \frac{1}{b-a} \int_{\frac{a+b}{2}}^b (t-b) [f'(t) - \lambda] dt. \end{aligned} \quad (2.7)$$

Remark 12. The identity (2.1) has many particular cases of interest.

If $x \in (a, b)$ is a point of differentiability for the absolutely continuous function $f : [a, b] \rightarrow \mathbb{C}$, then we have the equality:

$$\begin{aligned} f(x) + \left(\frac{a+b}{2} - x\right) f'(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ = \frac{1}{b-a} \int_a^x (t-a) [f'(t) - f'(x)] dt + \frac{1}{b-a} \int_x^b (t-b) [f'(t) - f'(x)] dt. \end{aligned} \quad (2.8)$$

In particular we have

$$\begin{aligned} f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \\ = \frac{1}{b-a} \int_a^{\frac{a+b}{2}} (t-a) \left[f'(t) - f'\left(\frac{a+b}{2}\right) \right] dt \\ + \frac{1}{b-a} \int_{\frac{a+b}{2}}^b (t-b) \left[f'(t) - f'\left(\frac{a+b}{2}\right) \right] dt \end{aligned} \quad (2.9)$$

provided $f'\left(\frac{a+b}{2}\right)$ exists and is finite.

For $x \in (a, b)$, if we take in (2.1)

$$\lambda_1(x) = \frac{f(x) - f(a)}{x-a} \quad \text{and} \quad \lambda_2(x) = \frac{f(b) - f(x)}{b-x},$$

then we get, after some elementary calculations,

$$\begin{aligned} & \frac{1}{2} \left[f(x) + \frac{(b-x)f(b) + (x-a)f(a)}{b-a} \right] - \frac{1}{b-a} \int_a^b f(t) dt \\ &= \frac{1}{b-a} \int_a^x (t-a) \left[f'(t) - \frac{f(x) - f(a)}{x-a} \right] dt \\ &+ \frac{1}{b-a} \int_x^b (t-b) \left[f'(t) - \frac{f(b) - f(x)}{b-x} \right] dt. \end{aligned} \quad (2.10)$$

In particular, we have

$$\begin{aligned} & \frac{1}{2} \left[f\left(\frac{a+b}{2}\right) + \frac{f(b) + f(a)}{2} \right] - \frac{1}{b-a} \int_a^b f(t) dt \\ &= \frac{1}{b-a} \int_a^{\frac{a+b}{2}} (t-a) \left[f'(t) - \frac{f\left(\frac{a+b}{2}\right) - f(a)}{\frac{b-a}{2}} \right] dt \\ &+ \frac{1}{b-a} \int_{\frac{a+b}{2}}^b (t-b) \left[f'(t) - \frac{f(b) - f\left(\frac{a+b}{2}\right)}{\frac{b-a}{2}} \right] dt. \end{aligned} \quad (2.11)$$

If we assume that the lateral derivatives $f'_+(a)$ and $f'_-(b)$ exist and are finite, then we have from (2.1) for $\lambda_1(x) = f'_+(a)$ and $\lambda_2(x) = f'_-(b)$

$$\begin{aligned} & f(x) + \frac{1}{2(b-a)} \left[(b-x)^2 f'_-(b) - (x-a)^2 f'_+(a) \right] - \frac{1}{b-a} \int_a^b f(t) dt \\ &= \frac{1}{b-a} \int_a^x (t-a) [f'(t) - f'_+(a)] dt \\ &+ \frac{1}{b-a} \int_x^b (t-b) [f'(t) - f'_-(b)] dt, \end{aligned} \quad (2.12)$$

for all $x \in [a, b]$.

In particular, we have

$$\begin{aligned} & f\left(\frac{a+b}{2}\right) + \frac{1}{8}(b-a) [f'_-(b) - f'_+(a)] - \frac{1}{b-a} \int_a^b f(t) dt \\ &= \frac{1}{b-a} \int_a^{\frac{a+b}{2}} (t-a) [f'(t) - f'_+(a)] dt \\ &+ \frac{1}{b-a} \int_{\frac{a+b}{2}}^b (t-b) [f'(t) - f'_-(b)] dt. \end{aligned} \quad (2.13)$$

If we take in (2.1) $\lambda_2(x) = \lambda_2(x) = f'\left(\frac{a+b}{2}\right)$, provided this derivative exists and is finite, then we get

$$\begin{aligned} & f(x) + \left(\frac{a+b}{2} - x\right) f'\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \\ &= \frac{1}{b-a} \int_a^x (t-a) \left[f'(t) - f'\left(\frac{a+b}{2}\right) \right] dt \\ &+ \frac{1}{b-a} \int_x^b (t-b) \left[f'(t) - f'\left(\frac{a+b}{2}\right) \right] dt, \end{aligned} \quad (2.14)$$

for all $x \in [a, b]$.

If we assume that the derivatives $f'_+(a)$, $f'_-(b)$ and $f'(x)$ exist and are finite, then by taking

$$\lambda_1(x) = \frac{f'_+(a) + f'(x)}{2} \text{ and } \lambda_2(x) = \frac{f'(x) + f'_-(b)}{2}$$

in (2.1) we get

$$\begin{aligned} & f(x) + \frac{1}{2} \left(\frac{a+b}{2} - x \right) f'(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ & + \frac{1}{4(b-a)} \left[(b-x)^2 f'_-(b) - (x-a)^2 f'_+(a) \right] \\ & = \frac{1}{b-a} \int_a^x (t-a) \left[f'(t) - \frac{f'_+(a) + f'(x)}{2} \right] dt \\ & + \frac{1}{b-a} \int_x^b (t-b) \left[f'(t) - \frac{f'(x) + f'_-(b)}{2} \right] dt. \end{aligned} \quad (2.15)$$

In particular, we have

$$\begin{aligned} & f\left(\frac{a+b}{2}\right) + \frac{1}{16} (b-a) [f'_-(b) - f'_+(a)] - \frac{1}{b-a} \int_a^b f(t) dt \\ & = \frac{1}{b-a} \int_a^{\frac{a+b}{2}} (t-a) \left[f'(t) - \frac{f'_+(a) + f'\left(\frac{a+b}{2}\right)}{2} \right] dt \\ & + \frac{1}{b-a} \int_{\frac{a+b}{2}}^b (t-b) \left[f'(t) - \frac{f'\left(\frac{a+b}{2}\right) + f'_-(b)}{2} \right] dt. \end{aligned} \quad (2.16)$$

3 Inequalities for Bounded Derivatives

Now, for $\gamma, \Gamma \in \mathbb{C}$ and $[a, b]$ an interval of real numbers, define the sets of complex-valued functions

$$\begin{aligned} & \bar{U}_{[a,b]}(\gamma, \Gamma) \\ & := \left\{ f : [a, b] \rightarrow \mathbb{C} \mid \operatorname{Re} \left[(\Gamma - f(t)) \left(\overline{f(t)} - \bar{\gamma} \right) \right] \geq 0 \text{ for almost every } t \in [a, b] \right\} \end{aligned}$$

and

$$\bar{\Delta}_{[a,b]}(\gamma, \Gamma) := \left\{ f : [a, b] \rightarrow \mathbb{C} \mid \left| f(t) - \frac{\gamma + \Gamma}{2} \right| \leq \frac{1}{2} |\Gamma - \gamma| \text{ for a.e. } t \in [a, b] \right\}.$$

The following representation result may be stated.

Proposition 13. *For any $\gamma, \Gamma \in \mathbb{C}$, $\gamma \neq \Gamma$, we have that $\bar{U}_{[a,b]}(\gamma, \Gamma)$ and $\bar{\Delta}_{[a,b]}(\gamma, \Gamma)$ are nonempty, convex and closed sets and*

$$\bar{U}_{[a,b]}(\gamma, \Gamma) = \bar{\Delta}_{[a,b]}(\gamma, \Gamma). \quad (3.1)$$

Proof. We observe that for any $z \in \mathbb{C}$ we have the equivalence

$$\left| z - \frac{\gamma + \Gamma}{2} \right| \leq \frac{1}{2} |\Gamma - \gamma|$$

if and only if

$$\operatorname{Re}[(\Gamma - z)(\bar{z} - \bar{\gamma})] \geq 0.$$

This follows by the equality

$$\frac{1}{4} |\Gamma - \gamma|^2 - \left| z - \frac{\gamma + \Gamma}{2} \right|^2 = \operatorname{Re}[(\Gamma - z)(\bar{z} - \bar{\gamma})]$$

that holds for any $z \in \mathbb{C}$.

The equality (3.1) is thus a simple consequence of this fact. \square

On making use of the complex numbers field properties we can also state that:

Corollary 14. *For any $\gamma, \Gamma \in \mathbb{C}$, $\gamma \neq \Gamma$, we have that*

$$\begin{aligned} \bar{U}_{[a,b]}(\gamma, \Gamma) = \{f : [a, b] \rightarrow \mathbb{C} \mid & (\operatorname{Re} \Gamma - \operatorname{Re} f(t))(\operatorname{Re} f(t) - \operatorname{Re} \gamma) \\ & + (\operatorname{Im} \Gamma - \operatorname{Im} f(t))(\operatorname{Im} f(t) - \operatorname{Im} \gamma) \geq 0 \text{ for a.e. } t \in [a, b]\}. \end{aligned} \quad (3.2)$$

Now, if we assume that $\operatorname{Re}(\Gamma) \geq \operatorname{Re}(\gamma)$ and $\operatorname{Im}(\Gamma) \geq \operatorname{Im}(\gamma)$, then we can define the following set of functions as well:

$$\begin{aligned} \bar{S}_{[a,b]}(\gamma, \Gamma) := \{f : [a, b] \rightarrow \mathbb{C} \mid & \operatorname{Re}(\Gamma) \geq \operatorname{Re} f(t) \geq \operatorname{Re}(\gamma) \\ & \text{and } \operatorname{Im}(\Gamma) \geq \operatorname{Im} f(t) \geq \operatorname{Im}(\gamma) \text{ for a.e. } t \in [a, b]\}. \end{aligned} \quad (3.3)$$

One can easily observe that $\bar{S}_{[a,b]}(\gamma, \Gamma)$ is closed, convex and

$$\emptyset \neq \bar{S}_{[a,b]}(\gamma, \Gamma) \subseteq \bar{U}_{[a,b]}(\gamma, \Gamma). \quad (3.4)$$

Theorem 15. *Let $f : [a, b] \rightarrow \mathbb{C}$ be an absolutely continuous on $[a, b]$ and $x \in (a, b)$. Suppose that $\gamma_i, \Gamma_i \in \mathbb{C}$ with $\gamma_i \neq \Gamma_i$, $i = 1, 2$ and $f' \in \bar{U}_{[a,x]}(\gamma_1, \Gamma_1) \cap \bar{U}_{[x,b]}(\gamma_2, \Gamma_2)$, then we have*

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right. \\ & + \frac{1}{2(b-a)} \left[(b-x)^2 \frac{\Gamma_2 + \gamma_2}{2} - (x-a)^2 \frac{\Gamma_1 + \gamma_1}{2} \right] \\ & \leq \frac{1}{4} \left[|\Gamma_1 - \gamma_1| \left(\frac{x-a}{b-a} \right)^2 + |\Gamma_2 - \gamma_2| \left(\frac{b-x}{b-a} \right)^2 \right] (b-a) \\ & \leq \frac{1}{4} (b-a) \\ & \times \begin{cases} \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] \max\{|\Gamma_1 - \gamma_1|, |\Gamma_2 - \gamma_2|\}. \\ \left[\left(\frac{x-a}{b-a} \right)^{2p} + \left(\frac{b-x}{b-a} \right)^{2p} \right]^{1/p} [|\Gamma_1 - \gamma_1|^q + |\Gamma_2 - \gamma_2|^q]^{1/q} \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ \left[\frac{1}{2} + \left| \frac{x - \frac{a+b}{2}}{b-a} \right| \right] [|\Gamma_1 - \gamma_1| + |\Gamma_2 - \gamma_2|]. \end{cases} \end{aligned} \quad (3.5)$$

Proof. Since $f' \in \bar{U}_{[a,x]}(\gamma_1, \Gamma_1) \cap \bar{U}_{[x,b]}(\gamma_2, \Gamma_2)$, then by taking the modulus in (2.1) for $\lambda_1(x) = \frac{\Gamma_1 + \gamma_1}{2}$ and $\lambda_2(x) = \frac{\Gamma_2 + \gamma_2}{2}$ we get

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right. \\ & \left. + \frac{1}{2(b-a)} \left[(b-x)^2 \frac{\Gamma_2 + \gamma_2}{2} - (x-a)^2 \frac{\Gamma_1 + \gamma_1}{2} \right] \right| \\ & \leq \frac{1}{b-a} \left| \int_a^x (t-a) \left[f'(t) - \frac{\Gamma_1 + \gamma_1}{2} \right] dt \right| \\ & \quad + \frac{1}{b-a} \left| \int_x^b (t-b) \left[f'(t) - \frac{\Gamma_2 + \gamma_2}{2} \right] dt \right| \\ & \leq \frac{1}{b-a} \int_a^x (t-a) \left| f'(t) - \frac{\Gamma_1 + \gamma_1}{2} \right| dt \\ & \quad + \frac{1}{b-a} \int_x^b (t-b) \left| f'(t) - \frac{\Gamma_2 + \gamma_2}{2} \right| dt \\ & \leq \frac{1}{b-a} \frac{|\Gamma_1 - \gamma_1|}{2} \int_a^x (t-a) dt + \frac{1}{b-a} \frac{|\Gamma_2 - \gamma_2|}{2} \int_x^b (b-t) dt \\ & = \frac{1}{4} \left[|\Gamma_1 - \gamma_1| \left(\frac{x-a}{b-a} \right)^2 + |\Gamma_2 - \gamma_2| \left(\frac{b-x}{b-a} \right)^2 \right] (b-a) \end{aligned}$$

and the first inequality in (3.5) is proved.

The last part follows by Hölder's inequality

$$mn + pq \leq (m^\alpha + p^\alpha)^{1/\alpha} (n^\beta + q^\beta)^{1/\beta},$$

where $m, n, p, q \geq 0$ and $\alpha > 1$ with $\frac{1}{\alpha} + \frac{1}{\beta} = 1$. □

Corollary 16. Let $f : [a, b] \rightarrow \mathbb{C}$ be an absolutely continuous on $[a, b]$ and $x \in (a, b)$. Suppose that $\gamma, \Gamma \in \mathbb{C}$ with $\gamma \neq \Gamma$, and $f' \in \bar{U}_{[a,b]}(\gamma, \Gamma)$, then we have

$$\begin{aligned} & \left| f(x) + \left(\frac{a+b}{2} - x \right) \frac{\Gamma + \gamma}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{2} |\Gamma - \gamma| \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] (b-a). \end{aligned} \quad (3.6)$$

In particular, we have

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{8} |\Gamma - \gamma| (b-a). \quad (3.7)$$

Remark 17. If the derivative $f' : [a, b] \rightarrow \mathbb{R}$ is bounded above and below, that is, there exists the constants $M > m$ such that

$$-\infty < m \leq f'(t) \leq M < \infty \text{ for a.e. } t \in [a, b],$$

then we recapture from (3.6) the inequality (1.6).

Remark 18. Let $f : [a, b] \rightarrow \mathbb{C}$ be an absolutely continuous on $[a, b]$. Suppose that $\gamma_i, \Gamma_i \in \mathbb{C}$ with $\gamma_i \neq \Gamma_i$, $i = 1, 2$ and $f' \in \bar{U}_{[a, \frac{a+b}{2}]}(\gamma_1, \Gamma_1) \cap \bar{U}_{[\frac{a+b}{2}, b]}(\gamma_2, \Gamma_2)$, then we have from (3.5) that

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt + \frac{1}{8}(b-a) \left(\frac{\Gamma_2 + \gamma_2}{2} - \frac{\Gamma_1 + \gamma_1}{2} \right) \right| \\ & \leq \frac{1}{16} [|\Gamma_1 - \gamma_1| + |\Gamma_2 - \gamma_2|] (b-a). \end{aligned} \quad (3.8)$$

4 Inequalities for Derivatives of Bounded Variation

Assume that the function $f : I \rightarrow \mathbb{C}$ is differentiable on the interior of I , denoted $\overset{\circ}{I}$, and $[a, b] \subset \overset{\circ}{I}$. Then, as in (2.15), we have the equality

$$\begin{aligned} & f(x) + \frac{1}{2} \left(\frac{a+b}{2} - x \right) f'(x) - \frac{1}{b-a} \int_a^b f(t) dt \\ & + \frac{1}{4(b-a)} \left[(b-x)^2 f'(b) - (x-a)^2 f'(a) \right] \\ & = \frac{1}{b-a} \int_a^x (t-a) \left[f'(t) - \frac{f'(a) + f'(x)}{2} \right] dt \\ & + \frac{1}{b-a} \int_x^b (t-b) \left[f'(t) - \frac{f'(x) + f'(b)}{2} \right] dt, \end{aligned} \quad (4.1)$$

for any $x \in [a, b]$.

Theorem 19. Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on $\overset{\circ}{I}$ and $[a, b] \subset \overset{\circ}{I}$. If the derivative $f' : \overset{\circ}{I} \rightarrow \mathbb{C}$ is of bounded variation on $[a, b]$, then

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt + \frac{1}{2} \left(\frac{a+b}{2} - x \right) f'(x) \right. \\ & \left. + \frac{1}{4(b-a)} \left[(b-x)^2 f'(b) - (x-a)^2 f'(a) \right] \right| \\ & \leq \frac{1}{4} \left[\left(\frac{x-a}{b-a} \right)^2 \bigvee_a^x(f') + \left(\frac{b-x}{b-a} \right)^2 \bigvee_x^b(f') \right] (b-a) \\ & \leq \frac{1}{4} (b-a) \\ & \times \begin{cases} \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] \left[\frac{1}{2} \bigvee_a^b(f') + \frac{1}{2} \left| \bigvee_a^x(f') - \bigvee_x^b(f') \right| \right], \\ \left[\left(\frac{x-a}{b-a} \right)^{2p} + \left(\frac{b-x}{b-a} \right)^{2p} \right]^{1/p} \left[\left[\bigvee_a^x(f') \right]^q + \left[\bigvee_x^b(f') \right]^q \right]^{1/q}, \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ \left[\frac{1}{2} + \left| \frac{x - \frac{a+b}{2}}{b-a} \right| \right] \bigvee_a^b(f'), \end{cases} \end{aligned} \quad (4.2)$$

for any $x \in [a, b]$.

Proof. Taking the modulus in (4.1) we have

$$\begin{aligned}
 & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt + \frac{1}{2} \left(\frac{a+b}{2} - x \right) f'(x) \right. \\
 & \quad \left. + \frac{1}{4(b-a)} \left[(b-x)^2 f'(b) - (x-a)^2 f'(a) \right] \right| \\
 & \leq \frac{1}{b-a} \left| \int_a^x (t-a) \left[f'(t) - \frac{f'(a) + f'(x)}{2} \right] dt \right| \\
 & \quad + \frac{1}{b-a} \left| \int_x^b (t-b) \left[f'(t) - \frac{f'(x) + f'(b)}{2} \right] dt \right| \\
 & \leq \frac{1}{b-a} \int_a^x (t-a) \left| f'(t) - \frac{f'(a) + f'(x)}{2} \right| dt \\
 & \quad + \frac{1}{b-a} \int_x^b (b-t) \left| f'(t) - \frac{f'(x) + f'(b)}{2} \right| dt.
 \end{aligned} \tag{4.3}$$

Since $f' : \dot{I} \rightarrow \mathbb{C}$ is of bounded variation on $[a, x]$ and $[x, b]$, then

$$\begin{aligned}
 \left| f'(t) - \frac{f'(a) + f'(x)}{2} \right| &= \frac{|f'(t) - f'(a) + f'(t) - f'(x)|}{2} \\
 &\leq \frac{1}{2} [|f'(t) - f'(a)| + |f'(x) - f'(t)|] \\
 &\leq \frac{1}{2} \bigvee_a^x(f')
 \end{aligned}$$

for any $t \in [a, x]$ and, similarly,

$$\left| f'(t) - \frac{f'(x) + f'(b)}{2} \right| \leq \frac{1}{2} \bigvee_x^b(f')$$

for any $t \in [x, b]$.

Then

$$\begin{aligned}
 \int_a^x (t-a) \left| f'(t) - \frac{f'(a) + f'(x)}{2} \right| dt &\leq \frac{1}{2} \bigvee_a^x(f') \int_a^x (t-a) dt \\
 &= \frac{1}{4} (x-a)^2 \bigvee_a^x(f')
 \end{aligned}$$

and

$$\begin{aligned}
 \int_x^b (b-t) \left| f'(t) - \frac{f'(x) + f'(b)}{2} \right| dt &\leq \frac{1}{2} \bigvee_x^b(f') \int_x^b (b-t) dt \\
 &= \frac{1}{4} (b-x)^2 \bigvee_x^b(f')
 \end{aligned}$$

and by (4.3) we get the desired inequality (4.2).

The last part follows by Hölder's inequality

$$mn + pq \leq (m^\alpha + p^\alpha)^{1/\alpha} (n^\beta + q^\beta)^{1/\beta},$$

where $m, n, p, q \geq 0$ and $\alpha > 1$ with $\frac{1}{\alpha} + \frac{1}{\beta} = 1$. □

Corollary 20. Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on $\overset{\circ}{I}$ and $[a, b] \subset \overset{\circ}{I}$. If the derivative $f' : \overset{\circ}{I} \rightarrow \mathbb{C}$ is of bounded variation on $[a, b]$, then

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt + \frac{1}{16} (b-a) [f'(b) - f'(a)] \right| \\ & \leq \frac{1}{16} (b-a) \bigvee_a^b(f'). \end{aligned} \quad (4.4)$$

Remark 21. If $p \in (a, b)$ is a median point in bounded variation for the derivative, i.e. $\bigvee_a^p(f') = \bigvee_p^b(f')$, then under the assumptions of Theorem 19, we have

$$\begin{aligned} & \left| f(p) - \frac{1}{b-a} \int_a^b f(t) dt + \frac{1}{2} \left(\frac{a+b}{2} - p\right) f'(p) \right. \\ & \quad \left. + \frac{1}{4(b-a)} [(b-p)^2 f'(b) - (p-a)^2 f'(a)] \right| \\ & \leq \frac{1}{8} (b-a) \left[\frac{1}{4} + \left(\frac{p - \frac{a+b}{2}}{b-a}\right)^2 \right] \bigvee_a^b(f'). \end{aligned} \quad (4.5)$$

5 Inequalities for Lipschitzian Derivatives

We say that $v : [a, b] \rightarrow \mathbb{C}$ is *Lipschitzian* with the constant $L > 0$, if it satisfies the condition

$$|v(t) - v(s)| \leq L|t - s| \text{ for any } t, s \in [a, b].$$

Theorem 22. Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on $\overset{\circ}{I}$ and $[a, b] \subset \overset{\circ}{I}$. Let $x \in (a, b)$. If the derivative $f' : \overset{\circ}{I} \rightarrow \mathbb{C}$ is Lipschitzian with the constant $K_1(x)$ on $[a, x]$ and constant $K_2(x)$ on $[x, b]$, then

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt + \frac{1}{2} \left(\frac{a+b}{2} - x\right) f'(x) \right. \\ & \quad \left. + \frac{1}{4(b-a)} [(b-x)^2 f'(b) - (x-a)^2 f'(a)] \right| \end{aligned} \quad (5.1)$$

$$\begin{aligned}
&\leq \frac{1}{8} \left[\left(\frac{x-a}{b-a} \right)^3 K_1(x) + \left(\frac{b-x}{b-a} \right)^3 K_2(x) \right] (b-a)^2 \\
&\leq \frac{1}{8} (b-a)^2 \\
&\quad \times \begin{cases} \left[\left(\frac{x-a}{b-a} \right)^3 + \left(\frac{b-x}{b-a} \right)^3 \right] \max \{ K_1(x), K_2(x) \}, \\ \left[\left(\frac{x-a}{b-a} \right)^{2p} + \left(\frac{b-x}{b-a} \right)^{2p} \right]^{1/p} [K_1^q(x) + K_2^q(x)]^{1/q} \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ \left[\frac{1}{2} + \left| \frac{x-\frac{a+b}{2}}{b-a} \right| \right]^3 [K_1(x) + K_2(x)]. \end{cases}
\end{aligned}$$

Proof. Since $f' : I \rightarrow \mathbb{C}$ is Lipschitzian with the constant $K_1(x)$ on $[a, x]$ and constant $K_2(x)$ on $[x, b]$, then

$$\begin{aligned}
\left| f'(t) - \frac{f'(a) + f'(x)}{2} \right| &= \frac{|f'(t) - f'(a) + f'(t) - f'(x)|}{2} \\
&\leq \frac{1}{2} [|f'(t) - f'(a)| + |f'(x) - f'(t)|] \\
&\leq \frac{1}{2} K_1(x) [|t-a| + |x-t|] \\
&= \frac{1}{2} K_1(x) (x-a)
\end{aligned}$$

for any $t \in [a, x]$ and, similarly,

$$\begin{aligned}
\left| f'(t) - \frac{f'(x) + f'(b)}{2} \right| &\leq \frac{1}{2} K_2(x) [|t-x| + |b-t|] \\
&= \frac{1}{2} K_2(x) (b-x)
\end{aligned}$$

for any $t \in [x, b]$.

Then

$$\begin{aligned}
\int_a^x (t-a) \left| f'(t) - \frac{f'(a) + f'(x)}{2} \right| dt &\leq \frac{1}{2} K_1(x) (x-a) \int_a^x (t-a) dt \\
&= \frac{1}{8} (x-a)^3 K_1(x)
\end{aligned}$$

and

$$\begin{aligned}
\int_x^b (b-t) \left| f'(t) - \frac{f'(x) + f'(b)}{2} \right| dt &\leq \frac{1}{2} K_2(x) (b-x) \int_x^b (b-t) dt \\
&= \frac{1}{8} (b-x)^3 K_2(x).
\end{aligned}$$

Making use of the inequality (4.3) we deduce the first bound in (5.1).

The second part is obvious. \square

Corollary 23. Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on \dot{I} and $[a, b] \subset \dot{I}$. If the derivative $f' : \dot{I} \rightarrow \mathbb{C}$ is Lipschitzian with the constant K on $[a, b]$ then

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt + \frac{1}{2} \left(\frac{a+b}{2} - x \right) f'(x) \right. \\ & \quad \left. + \frac{1}{4(b-a)} \left[(b-x)^2 f'(b) - (x-a)^2 f'(a) \right] \right| \\ & \leq \frac{1}{8} \left[\left(\frac{x-a}{b-a} \right)^3 + \left(\frac{b-x}{b-a} \right)^3 \right] K (b-a)^2 \end{aligned} \quad (5.2)$$

for any $x \in [a, b]$.

In particular, we have

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}\right) + \frac{1}{16}(b-a)[f'(b) - f'(a)] - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ & \leq \frac{1}{32} K (b-a)^2. \end{aligned} \quad (5.3)$$

References

- [1] A. M. Acu, A. Baboş and F. Sofonea, The mean value theorems and inequalities of Ostrowski type. *Sci. Stud. Res. Ser. Math. Inform.* **21** (2011), no. 1, 5–16.
- [2] A. M. Acu and F. Sofonea, On an inequality of Ostrowski type. *J. Sci. Arts* **2011**, no. 3(16), 281–287.
- [3] F. Ahmad, N. S. Barnett and S. S. Dragomir, New weighted Ostrowski and Čebyšev type inequalities. *Nonlinear Anal.* **71** (2009), no. 12, e1408–e1412.
- [4] M. W. Alomari, A companion of Ostrowski's inequality with applications. *Transylv. J. Math. Mech.* **3** (2011), no. 1, 9–14.
- [5] M. W. Alomari, M. Darus, S. S. Dragomir and P. Cerone, Ostrowski type inequalities for functions whose derivatives are s -convex in the second sense. *Appl. Math. Lett.* **23** (2010), no. 9, 1071–1076.
- [6] G. A. Anastassiou, Ostrowski type inequalities. *Proc. Amer. Math. Soc.* **123** (1995), No. 12, 3775–3781.
- [7] G. A. Anastassiou, Univariate Ostrowski inequalities, revisited. *Monatsh. Math.* **135** (2002), No. 3, 175–189.
- [8] G. A. Anastassiou, Ostrowski inequalities for cosine and sine operator functions. *Mat. Vesnik* **64** (2012), no. 4, 336–346.
- [9] G. A. Anastassiou, Multivariate right fractional Ostrowski inequalities. *J. Appl. Math. Inform.* **30** (2012), no. 3-4, 445–454.
- [10] G. A. Anastassiou, Univariate right fractional Ostrowski inequalities. *Cubo* **14** (2012), no. 1, 1–7.
- [11] N. S. Barnett, S. S. Dragomir and I. Gomm, A companion for the Ostrowski and the generalised trapezoid inequalities. *Math. Comput. Modelling* **50** (2009), no. 1-2, 179–187.
- [12] P. Cerone, W.-S. Cheung and S. S. Dragomir, On Ostrowski type inequalities for Stieltjes integrals with absolutely continuous integrands and integrators of bounded variation. *Comput. Math. Appl.* **54** (2007), No. 2, 183–191.
- [13] P. Cerone and S. S. Dragomir, Midpoint-type rules from an inequalities point of view. *Handbook of analytic-computational methods in applied mathematics*, 135–200, Chapman & Hall/CRC, Boca Raton, FL, 2000.

- [14] P. Cerone and S. S. Dragomir, Trapezoidal-type rules from an inequalities point of view. *Handbook of analytic-computational methods in applied mathematics*, 65–134, Chapman & Hall/CRC, Boca Raton, FL, 2000.
- [15] P. Cerone, S. S. Dragomir and C. E. M. Pearce, A generalised trapezoid inequality for functions of bounded variation, *Turk. J. Math.*, **24** (2000), 147–163.
- [16] S. S. Dragomir, The Ostrowski inequality for mappings of bounded variation, *Bull. Austral. Math. Soc.*, **60** (1999), 495–826.
- [17] S. S. Dragomir, On the mid-point quadrature formula for mappings with bounded variation and applications, *Kragujevac J. Math.*, **22** (2000), 13–19.
- [18] S. S. Dragomir, On the Ostrowski’s integral inequality for mappings with bounded variation and applications, *Math. Ineq. & Appl.*, **4**(1) (2001), 33–40. Preprint, *RGMIA Res. Rep. Coll.* **2**(1999), No. 1, Article 7. [Online: <http://rgmia.vu.edu.au/v2n1.html>].
- [19] S. S. Dragomir, On the trapezoid quadrature formula and applications, *Kragujevac J. Math.*, **23** (2001), 25–36.
- [20] S. S. Dragomir, Improvements of Ostrowski and generalised trapezoid inequality in terms of the upper and lower bounds of the first derivative. *Tamkang J. Math.* **34** (2003), no. 3, 213–222.
- [21] S. S. Dragomir, Refinements of the generalised trapezoid and Ostrowski inequalities for functions of bounded variation. *Arch. Math. (Basel)* **91** (2008), no. 5, 450–460.
- [22] S.S. Dragomir, Some inequalities for continuous functions of selfadjoint operators in Hilbert spaces, *Acta Math. Vietnamica*, to appear. Preprint *RGMIA Res. Rep. Coll.* **15**(2012), Art. 16. <http://rgmia.org/v15.php>.
- [23] S. S. Dragomir, Refinements of the Ostrowski inequality in terms of the cumulative variation and applications, Preprint *RGMIA Res. Rep. Coll.* **16** (2013), Art. 29, pp. 15 [Online: <http://rgmia.org/papers/v16/v16a29.pdf>].
- [24] Z. Liu, Some inequalities of perturbed trapezoid type. *J. Inequal. Pure Appl. Math.* **7** (2006), no. 2, Article 47, 9 pp.
- [25] Z. Liu, A note on Ostrowski type inequalities related to some s-convex functions in the second sense. *Bull. Korean Math. Soc.* **49** (2012), no. 4, 775–785.
- [26] Z. Liu, A sharp general Ostrowski type inequality. *Bull. Aust. Math. Soc.* **83** (2011), no. 2, 189–209.
- [27] Z. Liu, New sharp bound for a general Ostrowski type inequality. *Tamsui Oxf. J. Math. Sci.* **26** (2010), no. 1, 53–59.
- [28] Z. Liu, Some Ostrowski type inequalities and applications. *Vietnam J. Math.* **37** (2009), no. 1, 15–22.
- [29] Z. Liu, Some companions of an Ostrowski type inequality and applications. *J. Inequal. Pure Appl. Math.* **10** (2009), no. 2, Article 52, 12 pp.
- [30] M. Masjed-Jamei and S. S. Dragomir, A new generalization of the Ostrowski inequality and applications. *Filomat* **25** (2011), no. 1, 115–123.
- [31] B. G. Pachpatte, A note on a trapezoid type integral inequality. *Bull. Greek Math. Soc.* **49** (2004), 85–90.
- [32] J. Park, On the Ostrowskilike type integral inequalities for mappings whose second derivatives are s^* -convex. *Far East J. Math. Sci. (FJMS)* **67** (2012), no. 1, 21–35.
- [33] J. Park, Some Ostrowskilike type inequalities for differentiable real (α, m) -convex mappings. *Far East J. Math. Sci. (FJMS)* **61** (2012), no. 1, 75–91
- [34] M. Z. Sarikaya, On the Ostrowski type integral inequality. *Acta Math. Univ. Comenian. (N.S.)* **79** (2010), no. 1, 129–134.

- [35] W. T. Sulaiman, Some new Ostrowski type inequalities. *J. Appl. Funct. Anal.* **7** (2012), no. 1-2, 102–107.
- [36] K.-L. Tseng, Improvements of the Ostrowski integral inequality for mappings of bounded variation II. *Appl. Math. Comput.* **218** (2012), no. 10, 5841–5847.
- [37] K.-L. Tseng, S.-R. Hwang, G.-S. Yang, and Y.-M. Chou, Improvements of the Ostrowski integral inequality for mappings of bounded variation I. *Appl. Math. Comput.* **217** (2010), no. 6, 2348–2355.
- [38] N. Ujević, Error inequalities for a generalized trapezoid rule. *Appl. Math. Lett.* **19** (2006), no. 1, 32–37.
- [39] S. W. Vong, A note on some Ostrowskilike type inequalities. *Comput. Math. Appl.* **62** (2011), no. 1, 532–535.
- [40] Y. Wu and Y. Wang, On the optimal constants of Ostrowskilike inequalities involving n knots. *Appl. Math. Comput.* **219** (2013), no. 14, 7789–7794.
- [41] Q. Wu and S. Yang, A note to Ujević's generalization of Ostrowski's inequality. *Appl. Math. Lett.* **18** (2005), no. 6, 657–665.
- [42] Y.-X. Xiao, Remarks on Ostrowskilike inequalities. *Appl. Math. Comput.* **219** (2012), no. 3, 1158–1162.