

On Stability and Boundedness of Solutions of Certain Non Autonomous Fourth-Order Delay Differential Equations

Mebrouk Rahmane

Department of Mathematics, University of Oran 1, Algeria
mebroukrahmane@yahoo.fr

Moussadek Remili*

Department of Mathematics, University of Oran 1, Algeria
remilimous@gmail.com

Abstract

This paper establishes sufficient conditions to ensure the boundedness and asymptotic stability of solutions of a certain fourth order nonlinear non-autonomous differential equation by constructing Lyapunov functionals.

Received 24 February 2015

Revised 21 June 2015

Accepted in final form 17 April 2015

Communicated with Vladimír Janiš.

Keywords Stability, Lyapunov functional, fourth-order differential equations.

MSC(2010) 34C11.

1 Introduction

In applied science, some practical problems are associated with higher-order nonlinear differential equations, such as , electronic theory [22], biological model and other models [8] and [18] . Many results concerning the stability and boundedness of solutions of fourth order differential equations without delay have been obtained in view of various methods, especially, Lyapunov's method, see, the book of Reissig et al. [26] as a survey and the papers of Abou-El-Ela and Sadek [1], Adesina and Ogundare [3], Cartwright [9], Chukwu [10], Ezeilo [12], [14] Ezeilo and Tejumola [15], Harrow [16], Hu [17], Tejumola [30], Tunc [36], [37], [38], [39], Wu and Xiong [44] and the references cited therein. Besides, it should be noted that there are only a few results on the same problem for nonlinear differential equations of fourth order with delay, it have been discussed by a few authors, see, Bereketoglu [4], Abou-El-Ela et al. [2], Kang and Si [20], Sadek [27], Sinha [28], Tejumola [31], and Tunc [40], [41], [42]. The most efficient tool for the study of the stability and boundedness of solutions of a given nonlinear differential equation is provided by Lyapunov theory. But the construction of such functions which are positive definite with negative definite derivatives for higher-order differential equations with delay, is in general a difficult task.

*Corresponding Author

In [7], Chin has tried to use a new technique (the intrinsic method) to construct new Lyapunov functions for the following fourth-order differential equations

$$x'''' + a_1x'''' + a_2x'' + a_3x' + f(x) = 0, \tag{1.1}$$

$$x'''' + a_1x'''' + \psi(x')x'' + a_3x' + a_4x = 0, \tag{1.2}$$

$$x'''' + a_1x'''' + f(x, x')x'' + a_3x' + a_4x = 0, \tag{1.3}$$

in which a_1, a_2, a_3 and a_4 are constants. In [44], Wu and Xiong also investigated the asymptotic stability of the zero solution of the differential equations (1.1) and (1.3). Later, in 2004, Sadek [27] considered the fourth-order nonlinear delay differential equations of the form

$$x'''' + a_1x'''' + a_2x'' + a_3x' + f(x(t - r)) = 0,$$

and he derived sufficient conditions for the asymptotic stability of the zero-solution of these equations by constructing new Lyapunov functional.

In [33], Tunc investigated the asymptotic stability of zero solution of the fourth order non-linear differential equations with delay as follows

$$x'''' + \varphi(x'')x'''' + a_2x'' + a_3x' + f(x(t - r)) = 0.$$

In this article, we establish the uniform asymptotic stability of the differential equation of the form

$$\begin{aligned} \left(g(x(t))x''(t) \right)'' + a(t) \left(p(x(t))x''(t) \right)' + b(t) \left(q(x(t))x'(t) \right)' + c(t) f(x(t))x'(t) \\ + d(t) h(x(t - r)) = 0, \end{aligned} \tag{1.4}$$

where $g(x) > 0$ and r is a positive constant to be determined later; the primes in (1.4) denote differentiation with respect to t ; the functions a, b, c, d , are continuously differentiable functions. The functions f, g, h, p, q , are continuous. It is also supposed that the derivatives, $g'(x), p'(x), q'(x), f'(x)$ and $h'(x)$ exist and are continuous.

Equation (1.4) is equivalent to the system

$$\begin{cases} x' = y, \\ y' = \frac{1}{g(x)}z, \\ z' = w, \\ w' = -a(t)\frac{p(x)}{g(x)}w + \left(a(t)p(x)\theta_1(t) - b(t)\frac{q(x)}{g(x)} - a(t)g(x)\theta_2(t) \right) z \\ \quad - \left(b(t)g^2(x)\theta_3(t) + c(t)f(x) \right) y - d(t)h(x) + d(t) \int_{t-r}^t y(s)h'(x(s))ds, \end{cases} \tag{1.5}$$

where

$$\theta_1(t) = \frac{g'(x(t))}{g^2(x(t))}x'(t), \quad \theta_2(t) = \frac{p'(x(t))}{g^2(x(t))}x'(t), \quad \text{and} \quad \theta_3(t) = \frac{q'(x(t))}{g^2(x(t))}x'(t).$$

The continuity of the functions $a, b, c, d, f, g, g', h, p, p', q$, and q' guarantees the existence of the solutions of (1.4) (see [11], pp.15). It is assumed that the right hand side of the system (1.5) satisfies a Lipschitz condition in $x(t), y(t), z(t), w(t)$ and $x(t - r)$. This assumption guarantees the uniqueness of solutions of (1.4) ([11], pp.15). The motivation for the present paper comes from the results mentioned above. Our purpose is to

extend and improve the result established by Tunc [33], and Sadek [27] to the equation (1.4). Clearly the equation discussed in [27] and is a special case of equation (1.1) when $g(x) = p(x) = q(x) = 1$, and $a(t) = a$, $b(t) = b$, $c(t) = c$. Our approach is based on Lyapunov's second (direct) method. We shall use appropriate Lyapunov function and impose suitable conditions on the functions $g(x)$, $p(x)$, and $q(x)$.

2 Preliminaries

In this section, we shall state and prove certain results useful in the proof of our main result. Consider the functional differential equation

$$x' = f(t, x_t), \quad x_t(\theta) = x(t + \theta), \quad -r \leq \theta \leq 0, \quad t \geq 0, \quad (2.1)$$

where $f : I \times C_H \rightarrow \mathbb{R}^n$ is a continuous mapping, $f(t, 0) = 0$, $C_H := \{\phi \in (C[-r, 0], \mathbb{R}^n) : \|\phi\| \leq H\}$, and for $H_1 < H$, there exists $L(H_1) > 0$, with $|f(t, \phi)| < L(H_1)$ when $\|\phi\| < H_1$.

Definition 1. [6] An element $\psi \in C$ is in the ω -limit set of ϕ , say $\Omega(\phi)$, if $x(t, 0, \phi)$ is defined on $[0, +\infty)$ and there is a sequence $\{t_n\}$, $t_n \rightarrow \infty$, as $n \rightarrow \infty$, with $\|x_{t_n}(\phi) - \psi\| \rightarrow 0$ as $n \rightarrow \infty$ where $x_{t_n}(\phi) = x(t_n + \theta, 0, \phi)$ for $-r \leq \theta \leq 0$.

Definition 2. [6] A set $Q \subset C_H$ is an invariant set if for any $\phi \in Q$, the solution of (2.1), $x(t, 0, \phi)$, is defined on $[0, \infty)$ and $x_t(\phi) \in Q$ for $t \in [0, \infty)$.

Lemma 3. [5] If $\phi \in C_H$ is such that the solution $x_t(\phi)$ of (2.1) with $x_0(\phi) = \phi$ is defined on $[0, \infty)$ and $\|x_t(\phi)\| \leq H_1 < H$ for $t \in [0, \infty)$, then $\Omega(\phi)$ is a non-empty, compact, invariant set and

$$\text{dist}(x_t(\phi), \Omega(\phi)) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Lemma 4. [5] Let $V(t, \phi) : I \times C_H \rightarrow \mathbb{R}$ be a continuous functional satisfying a local Lipschitz condition, $V(t, 0) = 0$, and wedges W_i such that:

(i) $W_1(\|\phi\|) \leq V(t, \phi) \leq W_2(\|\phi\|)$.

(ii) $V'_{(2,1)}(t, \phi) \leq -W_3(\|\phi\|)$.

Then, the zero solution of (2.1) is uniformly asymptotically stable.

3 Assumptions and main results

First, we state some assumptions on the functions that appeared in (1.4), and suppose that there are positive constants $a_0, b_0, c_0, d_0, f_0, g_0, p_0, q_0, a_1, b_1, c_1, d_1, f_1, g_1, p_1, q_1, h_0, m, M, \delta, \delta_0, \eta_1$ and η_2 such that the following conditions are satisfied

i) $0 < a_0 \leq a(t) \leq a_1$; $0 < b_0 \leq b(t) \leq b_1$; $0 < c_0 \leq c(t) \leq c_1$;
 $0 < d_0 \leq d(t) \leq d_1$ for $t \geq 0$.

ii) $0 < f_0 \leq f(x) \leq f_1$; $g_0 \leq g(x) \leq g_1$; $0 < p_0 \leq p(x) \leq p_1$; $0 < q_0 \leq q(x) \leq q_1$
for $x \in \mathbb{R}$ and $0 < m < \min\{f_0, p_0, g_0, 1\}$, $M > \max\{f_1, g_1, p_1, 1\}$.

iii) $\frac{h(x)}{x} \geq \delta > 0$ (for $x \neq 0$); $h(0) = 0$.

iv) $\frac{h_0}{m} - \frac{a_0 m \delta_0}{d_1} \leq h'(x) \leq \frac{h_0}{2M}$ for $x \in \mathbb{R}$,

v) $b_0q_0 > \max(\kappa_1, \kappa_2)$ where

$$\left\{ \begin{aligned} \kappa_1 &= \frac{a_1h_0d_1M^2}{c_0m^3} + \frac{M^3(c_1 + \delta_0)}{a_0m^2} + a_0a_1m(M - 1), \\ \kappa_2 &= \frac{2d_1h_0a_0}{c_0(M - 1)} \left(\frac{1}{m} - \frac{1}{M} \right)^2 + 2\frac{c_0M}{a_0} + 2a_1\frac{d_1h_0M}{c_0m^3} + \frac{c_0c_1(M^2 + 2)mM}{d_1h_0}. \end{aligned} \right.$$

vi) $\int_0^{+\infty} (|a'(t)| + |b'(t)| + |c'(t)| + |d'(t)|) dt < \eta_1 < +\infty.$

vii) $\int_{-\infty}^{+\infty} (|g'(s)| + |p'(s)| + |q'(s)| + |f'(s)|) ds < \eta_2 < \infty.$

Now we dispose of the following lemma which will be required in the proof of next theorem.

Lemma 5. [19] Let $h(0) = 0, xh(x) > 0 (x \neq 0)$ and $\delta(t) - h'(x) \geq 0 (\delta(t) > 0)$, then

$$2\delta(t)H(x) \geq h^2(x) \quad \text{where} \quad H(x) = \int_0^x h(s)ds.$$

The main result of this paper is the following theorem.

Theorem 6. Suppose that assumptions i) ~ vii) hold. Then, every solution $x(t)$ of (1.4) and their derivatives $x'(t), x''(t)$ and $x'''(t)$ are uniformly asymptotically stable, provided that

$$r < \frac{1}{\lambda} \min \left\{ \epsilon c_0 m, \epsilon \frac{a_0 m}{M}, \frac{m^2(b_0 q_0 - \kappa_1) - \epsilon M^2(a_1 + c_1 m M)}{M m^2} \right\}, \tag{3.1}$$

where

$$\lambda = d_1 \lambda_0 (\alpha + \beta + 1), \quad \lambda_0 = \max \left\{ \frac{h_0}{2M}, \left| \frac{h_0}{m} - \frac{a_0 m \delta_0}{d_1} \right| \right\}, \quad \text{and} \tag{3.2}$$

$$\epsilon < \min \left\{ \frac{M}{a_0 m}, \frac{d_1 h_0}{c_0 m}, \frac{m^2(b_0 q_0 - \kappa_1)}{M^2(a_1 + m M c_1)} \right\}. \tag{3.3}$$

Proof. The proof depend on some fundamental properties of a continuously differentiable Lyapunov functional $W = W(t, x_t, y_t, z_t, w_t)$ defined by

$$W = e^{-\frac{1}{\eta} \int_0^t \gamma(s) ds} V, \tag{3.4}$$

where

$$\gamma(t) = |a'(t)| + |b'(t)| + |c'(t)| + |d'(t)| + |\theta_1(t)| + |\theta_2(t)| + |\theta_3(t)| + |\theta_4(t)|,$$

and

$$V = V_0(t, x_t, y_t, z_t, w_t) + \lambda \int_{-r}^0 \int_{t+s}^t y^2(\theta) d\theta ds,$$

such that

$$\begin{aligned}\theta_4(t) &= \frac{f'(x(t))}{g^2(x(t))} x'(t), \\ 2V_0 &= 2\beta d(t) H(x) + c(t) g(x) f(x) y^2 + \alpha b(t) \frac{q(x)}{g(x)} z^2 + a(t) \frac{p(x)}{g(x)} z^2 \\ &\quad + 2\beta a(t) \frac{p(x)}{g(x)} yz + [\beta b(t) q(x) - \alpha h_0 d(t)] y^2 - \beta \frac{1}{g(x)} z^2 + \alpha w^2 \\ &\quad + 2d(t) g(x) h(x) y + 2\alpha d(t) h(x) z + 2\alpha c(t) f(x) yz + 2\beta yw + 2zw,\end{aligned}$$

with $H(x) = \int_0^x h(s) ds$, $\alpha = \frac{M}{a_0 m} + \epsilon$, $\beta = \frac{d_1 h_0}{c_0 m} + \epsilon$ and η is positive constant to be determined later in the proof. $2V_0$ can be rearranged as the following

$$\begin{aligned}2V_0 &= a(t) p(x) \left(\frac{w}{a(t) p(x)} + z + \beta \frac{1}{g(x)} y \right)^2 + c(t) f(x) \left(\frac{d(t) h(x)}{c(t) f(x)} + y + \alpha z \right)^2 \\ &\quad + c(t) f(x) \left[\left(g(x) - 1 \right) y + \frac{d(t) h(x)}{c(t) f(x)} \right]^2 + 2\epsilon d(t) H(x) + V_1 + V_2 + V_3,\end{aligned}$$

where

$$\begin{aligned}V_1 &= 2d(t) \int_0^x h(s) \left(\frac{d_1 h_0}{c_0 m} - 2 \frac{d(t)}{c(t) f(x)} h'(s) \right) ds, \\ V_2 &= \left(\alpha b(t) \frac{q(x)}{g(x)} - \beta \frac{1}{g(x)} - \alpha^2 c(t) f(x) + a(t) p(x) \left(\frac{1}{g(x)} - 1 \right) \right) z^2,\end{aligned}$$

and

$$\begin{aligned}V_3 &= \left(\beta b(t) q(x) - \alpha h_0 d(t) - \beta^2 a(t) \frac{p(x)}{g^2(x)} - c(t) f(x) (g^2(x) - 3g(x) + 2) \right) y^2 \\ &\quad + \left(\alpha - \frac{1}{a(t) p(x)} \right) w^2 + 2\beta \left(1 - \frac{1}{g(x)} \right) yw.\end{aligned}$$

To prove that V is positive definite it suffice to show that V_1 , V_2 and V_3 are positive. Using conditions i) \sim v), and inequality (3.3) we obtain the following

$$\begin{aligned}V_1 &\geq 2d(t) \int_0^x 2h(s) \frac{d_1}{c_0 m} \left(\frac{h_0}{2} - h'(s) \right) ds \\ &\geq 4d_0 \frac{d_1}{c_0 m} \int_0^x h(s) \left(\frac{h_0}{2M} - h'(s) \right) ds \geq 0.\end{aligned}$$

Since (3.3) we get

$$\frac{M}{a_0 m} < \alpha < 2 \frac{M}{a_0 m}, \quad \frac{d_1 h_0}{c_0 m} < \beta < 2 \frac{d_1 h_0}{c_0 m}. \quad (3.5)$$

From (3.5) and rearrange V_2 we obtain

$$\begin{aligned}
 V_2 &= \alpha \left(b(t) \frac{g(x)}{g(x)} - \beta \frac{a(t)}{g(x)} - \alpha c(t) f(x) - \frac{a(t)p(x)}{\alpha} \left(1 - \frac{1}{g(x)} \right) \right) z^2 \\
 &\quad + \beta \left(\alpha \frac{a(t)}{g(x)} - \frac{1}{g(x)} \right) z^2 \\
 &\geq \alpha \left(\frac{b_0 q_0}{M} - \left(\frac{d_1 h_0}{c_0 m} + \epsilon \right) \frac{a_1}{m} - \left(\frac{M}{a_0 m} + \epsilon \right) c_1 M - a_1 \frac{a_0 m}{M} (M-1) \right) z^2 \\
 &\quad + \beta \left(\alpha \frac{a_0}{M} - \frac{1}{m} \right) z^2 \\
 &\geq \alpha \left(\frac{b_0 q_0}{M} - \frac{d_1 h_0 a_1}{c_0 m^2} - \frac{c_1 M^2}{a_0 m} - a_1 \frac{a_0 m}{M} (M-1) - \frac{\epsilon}{m} (a_1 + c_1 m M) \right) z^2 \\
 &\geq \frac{\alpha}{M m} \left(m(b_0 q_0 - \kappa_1) - \epsilon M (a_1 + c_1 m M) \right) z^2 \geq 0,
 \end{aligned}$$

and

$$\begin{aligned}
 V_3 &\geq \beta \left(b_0 q_0 - \frac{\alpha}{\beta} h_0 d_1 - a_1 \beta \frac{M}{g^2(x)} - \frac{c_1 M (M^2 + 2)}{\beta} \right) y^2 + \left(\frac{M-1}{a_0 m} \right) w^2 \\
 &\quad + 2\beta \left(1 - \frac{1}{g(x)} \right) yw \\
 &\geq \beta \left(b_0 q_0 - 2 \frac{M}{a_0} c_0 - 2a_1 \frac{d_1 h_0 M}{c_0 m^3} - \frac{c_0 c_1 (M^2 + 2) m M}{d_1 h_0} \right) y^2 + \left(\frac{M-1}{a_0 m} \right) w^2 \\
 &\quad + 2\beta \left(1 - \frac{1}{g(x)} \right) yw \\
 &\geq \psi(y, \omega),
 \end{aligned}$$

such that

$$\psi(y, \omega) = \beta \frac{2d_1 h_0 a_0}{c_0 (M-1)} \left(\frac{1}{m} - \frac{1}{M} \right)^2 y^2 + \left(\frac{M-1}{a_0 m} \right) w^2 + 2\beta \left(1 - \frac{1}{g(x)} \right) yw.$$

Observe that $\psi(y, \omega)$ is positive definite. Indeed by calculating the discriminant

$$\Delta = \beta^2 \left(1 - \frac{1}{g(x)} \right)^2 - \beta \frac{2d_1 h_0}{c_0 m} \left(\frac{1}{m} - \frac{1}{M} \right)^2,$$

since

$$\frac{1}{M} < \frac{1}{g(x)} < \frac{1}{m}, \text{ and } \frac{1}{M} < 1 < \frac{1}{m},$$

we get

$$\left| 1 - \frac{1}{g(x)} \right| < \frac{1}{m} - \frac{1}{M}.$$

Using (3.5) we obtain

$$\Delta \leq \beta \left[\frac{2d_1 h_0}{c_0 m} \left(\frac{1}{M} - \frac{1}{m} \right)^2 - \frac{2d_1 h_0}{c_0 m} \left(\frac{1}{M} - \frac{1}{m} \right)^2 \right] = 0.$$

Using the fact that the integral $\int_{-r}^0 \int_{t+s}^t y^2(\theta) d\theta ds$ is positive, we deduce that there exists positive number D_0 such that

$$2V \geq D_0 (y^2 + z^2 + w^2 + H(x)). \quad (3.6)$$

By lemma 5 and conditions iii) and iv) we conclude that there exists a positive number D_1 such that

$$2V \geq D_1 (x^2 + y^2 + z^2 + w^2), \quad (3.7)$$

thus V is positive definite which implies that W is also positive definite.

Then, we can find positive definite functions $U_1(\|X\|)$ and $U_2(\|X\|)$ such that $U_1(\|X\|) \leq V \leq U_2(\|X\|)$. By (ii) and (vii), we get

$$\begin{aligned} \int_0^t \left(\sum_{i=1}^4 |\theta_i(s)| \right) ds &= \int_{\alpha_1(t)}^{\alpha_2(t)} \frac{|g'(u)| + |p'(u)| + |q'(u)| + |f'(u)|}{g^2(u)} du \\ &\leq \frac{1}{m^2} \int_{-\infty}^{+\infty} (|g'(u)| + |p'(u)| + |q'(u)| + |f'(u)|) du < \infty, \end{aligned} \quad (3.8)$$

where $\alpha_1(t) = \min\{x(0), x(t)\}$, and $\alpha_2(t) = \max\{x(0), x(t)\}$. By inequalities (3.4), (3.7), and (3.8), we have

$$W \geq D_2(x^2 + y^2 + z^2 + w^2), \text{ where } D_2 = \frac{D_1}{2} e^{-\frac{1}{\eta}(\eta_1 + \frac{\eta_2}{m^2})}. \quad (3.9)$$

Therefore we can find positive definite functions $W_1(\|X\|)$ and $W_2(\|X\|)$ such that

$$W_1(\|X\|) \leq W \leq W_2(\|X\|).$$

Now we prove that \dot{W} is negative definite functional.

The derivative of V along any solution $(x(t), y(t), z(t), w(t))$ of system (1.5), we have

$$\begin{aligned} 2\dot{V}_{(1.5)} &= -2\epsilon c(t) f(x)y^2 + V_4 + V_5 + V_6 + V_7 + 2\frac{\partial V_0}{\partial t} + \lambda r y^2(t) - \lambda \int_{t-r}^t y^2(u) du \\ &\quad + 2\alpha w d(t) \int_{t-r}^t y(s) h'(x(s)) ds + 2\beta y d(t) \int_{t-r}^t y(s) h'(x(s)) ds \\ &\quad + 2z d(t) \int_{t-r}^t y(s) h'(x(s)) ds, \end{aligned}$$

where

$$\begin{aligned} V_4 &= -2 \left(\frac{d_1 h_0}{c_0 m} c(t) f(x) - d(t) g(x) h'(x) \right) y^2 - 2\alpha d(t) \left(\frac{h_0}{g(x)} - h'(x) \right) yz, \\ V_5 &= -2 \left(\frac{b(t) q(x)}{g(x)} - \alpha c(t) \frac{f(x)}{g(x)} - \beta a(t) \frac{p(x)}{g^2(x)} \right) z^2, \\ V_6 &= -2 \left(\alpha \frac{a(t) p(x)}{g(x)} - 1 \right) w^2 \end{aligned}$$

and

$$\begin{aligned} V_7 &= \theta_1 \left(a(t) p(x) z^2 - \alpha b(t) q(x) z^2 + c(t) f(x) g^2(x) y^2 + \beta z^2 + 2d(t) g^2(x) h(x) y \right. \\ &\quad \left. + 2\alpha a(t) p(x) z w \right) - b(t) \theta_3 g(x) \left(\alpha z^2 + 2\alpha g(x) z w + \beta g(x) y^2 + 2g(x) y z \right) \\ &\quad - a(t) \theta_2 g(x) \left(z^2 + 2\alpha z w \right) + \theta_4 \left(c(t) g^3(x) y^2 + 2\alpha c(t) g^2(x) y z \right). \end{aligned}$$

By conditions i), ii), iv), v) and inequalities (3.2), (3.3) and (3.5) we get

$$\begin{aligned}
 V_4 &\leq -2\left(d(t)h_0 - d(t)g(x)h'(x)\right)y^2 - 2\alpha d(t)\left(\frac{h_0}{g(x)} - h'(x)\right)yz \\
 &\leq -2d(t)g(x)\left(\frac{h_0}{g(x)} - h'(x)\right)y^2 - 2\alpha d(t)\left(\frac{h_0}{g(x)} - h'(x)\right)yz \\
 &\leq -2d(t)m\left(\frac{h_0}{g(x)} - h'(x)\right)\left[\left(y + \frac{\alpha}{2m}z\right)^2 - \left(\frac{\alpha}{2m}z\right)^2\right] \\
 &\leq -2d(t)m\left(\frac{h_0}{M} - h'(x)\right)\left(y + \frac{\alpha}{2m}z\right)^2 + 2d(t)m\left(\frac{h_0}{m} - h'(x)\right)\left(\frac{\alpha}{2m}z\right)^2 \\
 &\leq \frac{\alpha^2}{2m}d(t)\left(\frac{h_0}{m} - h'(x)\right)z^2.
 \end{aligned}$$

Hence,

$$\begin{aligned}
 V_4 + V_5 &\leq -2\left[b(t)\frac{g(x)}{g(x)} - \alpha c(t)\frac{f(x)}{g(x)} - \beta a(t)\frac{p(x)}{g^2(x)} - \frac{\alpha^2}{4m}d(t)\left(\frac{h_0}{m} - h'(x)\right)\right]z^2 \\
 &\leq -2\left[\frac{b_0q_0}{M} - \left(\frac{M}{a_0m} + \epsilon\right)\frac{c_1M}{m} - \left(\frac{d_1h_0}{c_0m} + \epsilon\right)\frac{a_1M}{m^2} - \frac{\alpha^2}{4m}(a_0\delta_0)\right]z^2 \\
 &\leq -2\left[\frac{b_0q_0}{M} - \frac{M^2}{a_0m^2}c_1 - \frac{d_1h_0a_1M}{c_0m^3} - \frac{M^2\delta_0}{a_0m^2} - \epsilon\frac{M}{m}\left(\frac{a_1}{m} + c_1\right)\right]z^2 \\
 &\leq -\frac{2}{Mm^2}\left(m^2(b_0q_0 - \kappa_1) - \epsilon M^2(a_1 + c_1m)\right)z^2 \leq 0.
 \end{aligned}$$

We have also,

$$V_6 \leq -2\left(\alpha\frac{a_0m}{M} - 1\right)w^2 = -2\epsilon\frac{a_0m}{M}w^2 \leq 0.$$

Putting $\lambda_1 = \min\left\{\epsilon c_0m, \epsilon\frac{a_0m}{M}, \frac{1}{Mm^2}\left(m^2(b_0q_0 - \kappa_1) - \epsilon M^2(a_1 + c_1mM)\right)\right\}$ and using Cauchy Schwartz inequality we have

$$\begin{aligned}
 &-2\epsilon c(t)f(x)y^2 + V_4 + V_5 + V_6 + 2\alpha wd(t)\int_{t-r}^t y(s)h'(x(s))ds - \lambda\int_{t-r}^t y^2(s)ds \\
 &+ 2\beta yd(t)\int_{t-r}^t y(s)h'(x(s))ds + 2zd(t)\int_{t-r}^t y(s)h'(x(s))ds + \lambda ry^2(t) \\
 &\leq -2\lambda_1(y^2 + z^2 + w^2) + \alpha d_1\lambda_0\left(w^2r + \int_{t-r}^t y^2(s)ds\right) + \beta d_1\lambda_0\left(y^2r + \int_{t-r}^t y^2(s)ds\right) \\
 &\quad + d_1\lambda_0\left(z^2r + \int_{t-r}^t y^2(s)ds\right) + \lambda ry^2(t) - \lambda\int_{t-r}^t y^2(s)ds \\
 &\leq -2\lambda_1(y^2 + z^2 + w^2) + d_1\lambda_0r(\beta y^2 + z^2 + \alpha w^2) + (d_1\lambda_0(\alpha + \beta + 1) - \lambda)\int_{t-r}^t y^2(s)ds \\
 &\leq -2\lambda_1(y^2 + z^2 + w^2) + d_1\lambda_0(\alpha + \beta + 1)r(y^2 + z^2 + w^2) \\
 &\leq -2D_3(y^2 + z^2 + w^2). \tag{3.10}
 \end{aligned}$$

Where $D_3 = \lambda_1 - \lambda r$. It can be seen that if $r < \frac{\lambda_1}{\lambda}$, then $D_2 > 0$.

Now by the inequalities (3.6), (3.10), the lemma 5 and the Cauchy Schwartz inequality

we get the following.

$$\begin{aligned}
 V_7 &\leq |\theta_1| \left(a(t) p(x) z^2 + \alpha b(t) q(x) z^2 + c(t) f(x) g^2(x) y^2 + \beta z^2 + d(t) g^2(x) (h^2(x) + y^2) \right. \\
 &\quad \left. + \alpha a(t) p(x) (z^2 + w^2) \right) + |\theta_4| \left(c(t) g^3(x) y^2 + \alpha c(t) g^2(x) (y^2 + z^2) \right) \\
 &\quad + b(t) |\theta_3| g(x) \left(\alpha z^2 + \alpha g(x) (z^2 + w^2) + \beta g(x) y^2 + g(x) (y^2 + z^2) \right) \\
 &\quad + a(t) |\theta_2| g(x) \left(z^2 + \alpha (z^2 + w^2) \right) \\
 &\leq \lambda_2 (|\theta_1| + |\theta_2| + |\theta_3| + |\theta_4|) (y^2 + z^2 + w^2 + H(x)) \\
 &\leq \frac{2\lambda_2}{D_0} (|\theta_1| + |\theta_2| + |\theta_3| + |\theta_4|) V.
 \end{aligned}$$

such that,

$$\lambda_2 = \max \left\{ d_1 h_0 M, \alpha M (a_1 + M b_1), c_1 M^3 + (d_1 + \alpha c_1 + \beta b_1 + b_1) M^2, \right. \\
 \left. \beta + a_1 M (\alpha + 1) + b_1 M (2\alpha + 1) + \alpha c_1 M^2 \right\}.$$

Similarly we have

$$\begin{aligned}
 2 \frac{\partial V_0}{\partial t} &= d'(t) [2\beta H(x) - \alpha h_0 y^2 + 2g(x) h(x) y + 2\alpha h(x) z] \\
 &\quad + c'(t) [g(x) f(x) y^2 + 2\alpha f(x) y z] + b'(t) \left[\alpha \frac{q(x)}{g(x)} z^2 + \beta q(x) y^2 \right] \\
 &\quad + a'(t) \left[\frac{p(x)}{g(x)} z^2 + 2\beta \frac{p(x)}{g(x)} y z \right].
 \end{aligned}$$

There exist positive constant λ_3 such that

$$\begin{aligned}
 2 \left| \frac{\partial V_0}{\partial t} \right| &\leq |d'(t)| \left(2\beta H(x) + \alpha h_0 y^2 + g(x) (h^2(x) + y^2) + \alpha (h^2(x) + z^2) \right) \\
 &\quad + |c'(t)| |f(x)| \left(g(x) y^2 + \alpha (y^2 + z^2) \right) + |b'(t)| |q(x)| \left(\alpha \frac{1}{g(x)} z^2 + \beta y^2 \right) \\
 &\quad + |a'(t)| \left| \frac{p(x)}{g(x)} \right| \left(z^2 + \beta (y^2 + z^2) \right) \\
 &\leq \lambda_3 \left(|a'(t)| + |b'(t)| + |c'(t)| + |d'(t)| \right) (y^2 + z^2 + w^2 + H(x)) \\
 &\leq 2 \frac{\lambda_3}{D_0} \left(|a'(t)| + |b'(t)| + |c'(t)| + |d'(t)| \right) V,
 \end{aligned}$$

where

$$\lambda_3 = \max \left\{ 2\beta + h_0 (\alpha + 1), M^2 + \beta \frac{M}{m} + \alpha (h_0 + M), \frac{M}{m} (\alpha + \beta + 1) \right\}.$$

Thus for $\frac{1}{\eta} = \frac{1}{D_0} \max \{ \lambda_2, \lambda_3 \}$ we have

$$\begin{aligned}
 \dot{V}_{(1.5)} &\leq -D_3 (y^2 + z^2 + w^2) + \frac{1}{\eta} \left(|a'(t)| + |b'(t)| + |c'(t)| + |d'(t)| \right. \\
 &\quad \left. + |\theta_1| + |\theta_2| + |\theta_3| + |\theta_4| \right) V.
 \end{aligned} \tag{3.11}$$

By conditions vi), vii) and inequalities (3.8), (3.11) we have

$$\begin{aligned} \dot{W}_{(1.5)} &= \left(\dot{V}_{(1.5)} - \frac{1}{\eta} \gamma(t) V \right) e^{-\frac{1}{\eta} \int_0^t \gamma(s) ds} \\ &\leq -D_3 (y^2 + z^2 + w^2) e^{-\frac{1}{\eta} \int_0^t \gamma(s) ds} \\ &\leq -D_4 (y^2 + z^2 + w^2), \end{aligned}$$

where $D_4 = D_3 e^{-\frac{\eta_1 + \eta_2}{\eta}}$. From (1.5), $W_3(\|X\|) = D_4(y^2 + z^2 + w^2)$ is positive definite function. Thus, we conclude that the solutions of system (1.5) are uniformly asymptotically stable. Now, it is evident from (1.5) that

$$\begin{cases} |x'(t)| = |y(t)|, \\ |x''(t)| = \left| \frac{z(t)}{g(x)} \right| \leq \frac{|z(t)|}{m}, \\ |x'''(t)| = \left| \frac{w(t)}{g(x)} - \frac{g'(x)x'(t)}{g^2(x)} \right| \leq \frac{|w(t)|}{m} + \frac{|g'(x)||x'(t)|}{m^2}. \end{cases}$$

Clearly, from the above discussion

$$\lim_{t \rightarrow \infty} x(t) = 0, \quad \lim_{t \rightarrow \infty} x'(t) = 0, \quad \lim_{t \rightarrow \infty} x''(t) = 0, \quad \text{and} \quad \lim_{t \rightarrow \infty} x'''(t) = 0.$$

This fact completes the proof of the Theorem. □

4 Example

We consider the following fourth order non-autonomous delay differential equation

$$\begin{aligned} &\left(\left(\frac{x^2(t) \sin x(t) + 5x^4(t) + 5}{5(1+x^4(t))} \right) x''(t) \right)'' \\ &+ (e^{-t} \sin t + 2) \left(\left(\frac{x(t) + 4e^{x(t)} + 4e^{-x(t)}}{4(e^{x(t)} + e^{-x(t)})} \right) x''(t) \right)' \\ &+ \left(\frac{\cos t + 7t^2 + 7}{1+t^2} \right) \left(\left(\frac{\sin x(t) + 6e^{x(t)} + 6e^{-x(t)}}{e^{x(t)} + e^{-x(t)}} \right) x'(t) \right)' \\ &+ (e^{-2t} \sin^3 t + 2) \left(\frac{x(t) \cos x(t) + 5x^4(t) + 5}{5(1+x^4(t))} \right) x'(t) \\ &+ \left(\frac{\cos^2 t + t^2 + 1}{10(1+t^2)} \right) \left(\frac{x(t-r)}{x^2(t-r) + 1} \right) = 0. \end{aligned} \tag{4.1}$$

By taking $g(x) = \frac{x^2 \sin x + 5x^4 + 5}{5(1+x^4)}$, $p(x) = \frac{x + 4e^x + 4e^{-x}}{4(e^x + e^{-x})}$, $q(x) = \frac{\sin x + 6e^x + 6e^{-x}}{e^x + e^{-x}}$,
 $f(x) = \frac{x \cos x + 5x^4 + 5}{5(1+x^4)}$, $h(x) = \frac{x}{x^2 + 1}$, $a(t) = e^{-t} \sin t + 2$, $b(t) = \frac{\cos t + 7t^2 + 7}{1+t^2}$,
 $c(t) = e^{-2t} \sin^3 t + 2$ and $d(t) = \frac{\cos^2 t + t^2 + 1}{10(1+t^2)}$.

We have

$$m = \frac{9}{10}, \quad M = \frac{11}{10}, \quad q_0 = \frac{11}{2}, \quad q_1 = \frac{13}{2}, \quad h_0 = \frac{5}{2}, \quad \delta_0 = \frac{5}{3}, \quad a_0 = 1, \quad a_1 = 3, \\ b_0 = 6, \quad b_1 = 8, \quad c_0 = 1, \quad c_1 = 3, \quad d_0 = \frac{1}{10}, \quad d_1 = \frac{1}{5}, \quad \text{and}$$

$$\frac{h_0}{m} - \frac{a_0 m \delta_0}{d_1} \leq -4.55 \leq h'(x) \leq 1.1 \leq \frac{h_0}{2M}, \\ \kappa_1 = \frac{a_1 h_0 d_1 M^2}{c_0 m^3} + \frac{M^3(c_1 + \delta_0)}{a_0 m^2} + a_0 a_1 m (M - 1) \leq 11, \\ \kappa_2 = \frac{2d_1 h_0 a_0}{c_0 (M - 1)} \left(\frac{1}{m} - \frac{1}{M} \right)^2 + 2 \frac{c_0 M}{a_0} + 2a_1 \frac{d_1 h_0 M}{c_0 m^3} + \frac{c_0 c_1 (M^2 + 2)mM}{d_1 h_0} \\ \leq 27, \\ \epsilon < \min \left\{ \frac{M}{a_0 m}, \frac{d_1 h_0}{c_0 m}, \frac{m^2(b_0 q_0 - \kappa_1)}{M^2(a_1 + mM c_1)} \right\} = \frac{5}{9}.$$

By choosing $\epsilon = \frac{1}{4}$ we get $\alpha = \frac{M}{a_0 m} + \epsilon = \frac{53}{36}, \quad \beta = \frac{d_1 h_0}{c_0 m} + \epsilon = \frac{29}{36},$

$$\lambda_0 = \max \left\{ \frac{h_0}{2M}, \left| \frac{h_0}{m} - \frac{a_0 m \delta_0}{d_1} \right| \right\} = \frac{85}{18}, \\ \lambda = d_1 \lambda_0 (\alpha + \beta + 1) = \frac{1003}{324}, \\ r < \frac{1}{\lambda} \min \left\{ \epsilon c_0 m, \epsilon \frac{a_0 m}{M}, \frac{m^2(b_0 q_0 - \kappa_1) - \epsilon M^2(a_1 + c_1 m M)}{M m^2} \right\} = \frac{729}{11033}.$$

On the other hand,

$$\int_{-\infty}^{+\infty} |g'(x)| dx = \frac{1}{5} \int_{-\infty}^{+\infty} \left| \frac{-4x^5 \sin x + (2x \sin x + x^2 \cos x)(x^4 + 1)}{(x^4 + 1)^2} \right| dx \leq \frac{3}{5} \sqrt{2}\pi, \\ \int_{-\infty}^{+\infty} |p'(x)| dx = \frac{1}{4} \int_{-\infty}^{+\infty} \left| \frac{1}{e^x + e^{-x}} + x \frac{e^{-x} - e^x}{(e^x + e^{-x})^2} \right| dx \leq \frac{\pi}{4},$$

$$\int_{-\infty}^{+\infty} |q'(x)| dx = \frac{1}{5} \int_{-\infty}^{+\infty} \left| \frac{(e^x + e^{-x}) \cos x - (e^x - e^{-x}) \sin x}{(e^x + e^{-x})^2} \right| dx \leq \frac{\pi}{5}, \\ \int_{-\infty}^{+\infty} |f'(x)| dx = \frac{1}{5} \int_{-\infty}^{+\infty} \left| \frac{(\cos x - x \sin x)(x^4 + 1) - 4x^4 \cos x}{(x^4 + 1)^2} \right| dx \\ \leq \frac{1}{5} \int_{-\infty}^{+\infty} \frac{5 + x^2}{x^4 + 1} dx = \frac{9}{10} \sqrt{2}\pi,$$

then,

$$\int_{-\infty}^{+\infty} (|g'(s)| + |p'(s)| + |q'(s)| + |f'(s)|) ds < \infty.$$

We have also

$$\begin{aligned} \int_0^{+\infty} |a'(t)| dt &= \int_0^{+\infty} |(\cos t)e^{-t} - (\sin t)e^{-t}| dt \leq \int_0^{+\infty} 2e^{-t} dt = 2, \\ \int_0^{+\infty} |b'(t)| dt &= \int_0^{+\infty} \left| -\frac{\sin t}{t^2+1} - 2t \frac{\cos t}{(t^2+1)^2} \right| dt \leq \int_0^{+\infty} \left(\frac{1}{t^2+1} + \frac{2t}{(t^2+1)^2} \right) dt \\ &\leq \int_0^{+\infty} \frac{2}{t^2+1} dt = \pi, \\ \int_0^{+\infty} |c'(t)| dt &= \int_0^{+\infty} |3(\cos t \sin^2 t)e^{-2t} - 2(\sin^3 t)e^{-2t}| dt \leq \int_0^{+\infty} 5e^{-2t} dt = \frac{5}{2}, \\ \int_0^{+\infty} |d'(t)| dt &= \int_0^{+\infty} \left| -2(\cos t) \frac{\sin t}{t^2+1} - 2t \frac{\cos^2 t}{(t^2+1)^2} \right| dt \leq \int_0^{+\infty} \frac{3}{t^2+1} dt = \frac{3\pi}{2}. \end{aligned}$$

Hence

$$\int_0^{+\infty} (|a'(t)| + |b'(t)| + |c'(t)| + |d'(t)|) dt < +\infty.$$

Thus all the assumptions of Theorem (6) hold, this shows that every solution $x(t)$ of (4.1) and their derivatives $x'(t)$, $x''(t)$ and $x'''(t)$ are uniformly asymptotically stable.

References

- [1] A. M. A. Abou-El-Ela, A. I. Sadek, A stability result for certain fourth order differential equations. *Ann. Differential Equations* 6 (1990), no. 1, 1–9.
- [2] A. M. A. Abou-El-Ela, A. I. Sadek, A. M. Mahmoud, On the stability of solutions of certain fourth-order nonlinear nonautonomous delay differential equation, *Int. J. Appl. Math*, 22 (2009), no. 2, 245–258.
- [3] O. A. Adesina, B. S. Ogundare, Some new stability and boundedness results on a certain fourth order nonlinear differential equation, *Nonlinear Stud*, 19 (2012), no. 3, 359–369.
- [4] H. Bereketoglu, Asymptotic stability in a fourth order delay differential equation, *Dynam. Systems Appl*, 7 (1998), no. 1, 105–115.
- [5] T. A. Burton, Stability and periodic solutions of ordinary and functional differential equations, *Mathematics in science and engineering*, Volume 178, Academic Press, INC, 1985.
- [6] T.A.Burton, Volterra Integral and Differential Equations, *Mathematics in Science and Engineering* V(202)(2005), 2nd edition.
- [7] P. S. M. Chin, Stability results for the solutions of certain fourth-order autonomous differential equations, *Internat. J. Control*, 49 (1989), no. 4, 1163–1173.
- [8] J. Cronin, Some mathematics of biological oscillations, *Siam Rev*, 19 (1977), 100–137.
- [9] M. L. Cartwright, On the stability of solutions of certain differential equations of the fourth order, *Quart. J. Mech. Appl. Math*, 9 (1956), 185–194.
- [10] E. N. Chukwu, On the stability of a nonhomogeneous differential equation of the fourth order, *Ann. Mat. Pura Appl*, (4) 92 (1972), 1–11.
- [11] L. Elsgolts, Introduction to the theory of differential equations with deviating arguments, *Translated from the Russian by Robert J. McLaughlin Holden-Day, Inc., San Francisco, Calif.-London-Amsterdam*, 1966.
- [12] J. O. C. Ezeilo, A stability result for solutions of a certain fourth order differential equation, *J. London Math. Soc*, 37 (1962), 28–32.

- [13] J. O. C. Ezeilo, On the boundedness and the stability of solutions of some differential equations of the fourth order, *J. Math. Anal. Appl.* 5 (1962), 136–146.
- [14] J. O. C. Ezeilo, Stability results for the solutions of some third and fourth order differential equations. *Ann. Mat. Pura Appl.* (4) 66 (1964), 233–249.
- [15] J. O. C. Ezeilo, H. O. Tejumola, On the boundedness and the stability properties of solutions of certain fourth order differential equations. *Ann. Mat. Pura Appl.* (4) 95 (1973), 131–145.
- [16] M. Harrow, Further results on the boundedness and the stability of solutions of some differential equations of the fourth order, *SIAM J. Math. Anal.* 1(1970), 189–194.
- [17] Hu. Chao Yang, The stability in the large for certain fourth order differential equations, *Ann. Differential Equations*, 8 (1992), no. 4, 422–428.
- [18] K. Gopalsamy, Stability and Oscillations in Delay Differential Equations of Population Dynamic, *Mathematics and Its Applications*, vol. 74, Kluwer Academic, Dordrecht, 1992.
- [19] T. Hara, On the asymptotic behavior of the solutions of some third and fourth order non-autonomous differential equations, *Publ. RIMS, Kyoto Univ.* 9 (1974), 649–673.
- [20] Huiyan. Kang, Si. Ligeng, Stability of solutions to certain fourth order delay differential equations, *Ann. Differential Equations*, 26 (2010), no. 4, 407–413.
- [21] P. S. M. Omeike, Boundedness of solutions of the fourth order differential equations with oscillatory restoring and forcing terms, *Analele stiintifice ale univesitatii "AL.I. CUZA" DIN IASI (S.N.) Matematica, Tomul LIV*, 2008, f.1
- [22] L.L. Rauch, Oscillations of a third order nonlinear autonomous system, *Contributions to the Theory of Nonlinear Oscillations*, *Ann. Math. Stud* 20 (1950), 39–88.
- [23] M. Remili, D. L. Oudjedi, Uniform stability and boundedness of a kind of third order delay differential equations, *Bull. Comput. Appl. Math.*, Vol 2, No1, 2014.
- [24] M. Remili, D. L. Oudjedi, Stability and boundedness of the solutions of non autonomous third order differential equations with delay, *cta Univ. Palacki. Olomuc., Fac. rer. nat, Mathematica*, Vol 53, No2, 2014.
- [25] M. Remili, D. Beldjerd, On the asymptotic behavior of the solutions of third order delay differential equations, *Rend. Circ. Mat. Palermo*, Vol 63, No 3, 2014.
- [26] R. Reissig, G. Sansone, R. Conti, Non-linear differential equations of higher order, *Translated from the German. Noordhoff International Publishing, Leyden*, 1974.
- [27] A. I. Sadek, On the stability of solutions of certain fourth order delay differential equations, *Applied Mathematics and Computation*, 148(2), (2004), 587–597.
- [28] A. S. C. Sinha, On stability of solutions of some third and fourth order delay-differential equations. *Information and Control* 23 (1973), 165–172.
- [29] Shair A.; *Asymptotic properties of linear fourth order differential equations*. American mathematical society volume 59 (1976), number 1, august.
- [30] H. O. Tejumola, Further results on the boundedness and the stability of certain fourth order differential equations, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* (8) 52 (1972), 16–23.
- [31] Tejumola, H.O.; Tchegnani, B., Stability, boundedness and existence of periodic solutions of some third and fourth order nonlinear delay differential equations, *J. Nigerian Math. Soc.* 19 (2000), 9–19.
- [32] C. Tunc and A. Tiryaki, Constructing Liapunov functions for some fourth-order autonomous differential equations. *Indian J pure applied math.* 26(3) March (1995), 225–232.
- [33] C. Tunc, Some stability results for the solutions of certain fourth order delay differential equations, *Differential Equations and Applications*, 4 (2004), 165–174.

- [34] C.Tunc, Stability and boundedness of solutions to certain fourth order differential equations. *Electronic Journal of Differential Equations*, Vol. 2006(2006), No. 35, pp. 1–10.
- [35] C.Tunc, Some remarks on the stability and boundedness of solutions of certain differential equations of fourth-order. *Computational and Applied Mathematics*, Volume 26 (2007), N. 1, pp. 1–17.
- [36] C.Tunc, On the stability and the boundedness properties of solutions of certain fourth order differential equations. *Istanbul. Univ. Fen Fak. Mat. Derg*, 54 (1995), 161–173.
- [37] C.Tunc, A note on the stability and boundedness results of solutions of certain fourth order differential equations. *Appl. Math. Comput*, 155 (2004), no. 3, 837–843.
- [38] C.Tunc, Some stability and boundedness results for the solutions of certain fourth order differential equations, *Acta Univ. Palack. Olomuc. Fac. Rerum Natur*, Math. 44 (2005), 161–171.
- [39] C.Tunc, Stability and boundedness of solutions to certain fourth order differential equations, *Electron. J. Differential Equations* (2006), No. 35, 10 pp.
- [40] C.Tunc, On the stability of solutions of non-autonomous differential equations of fourth order with delay, *Funct. Differ. Equ*, 17 (2010), no. 1-2, 195–212.
- [41] C.Tunc, On the stability and boundedness of solutions in a class of nonlinear differential equations of fourth order with constant delay, *Vietnam J. Math*, 38 (2010), no. 4, 453–466.
- [42] C.Tunc, The boundedness to nonlinear differential equations of fourth order with delay, *Nonlinear Stud.* 17(2010),no.1, 47–56.
- [43] V. Vlček, On the boundedness of solutions of a certain fourth-order nonlinear differential equation, *Acta Universitatis Palackianae Olomucensis. Facultas Rerum Naturalium. Mathematica*, Vol. 27 (1988), No. 1, 273–288.
- [44] X.Wu, K.Xiong, Remarks on stability results for the solutions of certain fourth-order autonomous differential equations, *Internat. J. Control.* 69, 2 (1998), 353–360.