

# Coincidence point theorems for a family of multivalued mappings in partially ordered metric spaces

**Binayak S. Choudhury**

*Department of Mathematics, Bengal Engineering and Science University, Shibpur, Howrah - 711103,  
West Bengal, India*

`binayak12@yahoo.co.in, binayak@becs.ac.in`

**Nikhilesh Metiya\***

*Department of Mathematics, Bengal Institute of Technology, Kolkata - 700150, West Bengal, India*

`metiya.nikhilesh@gmail.com`

---

## Abstract

In this paper we establish certain multivalued coincidence point results of a family of multivalued mappings with a singlevalued mapping under the assumptions of certain almost contractive type inequalities. Our results are derived in metric spaces with a partial ordering. The corresponding singlevalued cases are shown to extend a number of existing results. We have given one illustrative example. The methodology applied here is a blending of order theoretic and analytic methodologies.

*Received June 29, 2012*

*Revised January 25, 2013*

*Accepted in final form January 30, 2013*

*Communicated with Lubomír Snoha.*

**Keywords** Partial ordering, Metric space, Almost contraction, Coincidence point.

**MSC(2010)** 54H10, 54H25, 47H10.

---

## 1 Introduction

In the fixed point theory of setvalued maps two types of distances are generally used. One is the Hausdorff distance. Nadler [22] had proved a multivalued version of the Banach's contraction mapping principle by using the Hausdorff metric. There are many other fixed point results using this Hausdorff metric, some instances of these works are in [9, 17, 29, 30, 31]. The another distance is the  $\delta$  - distance. This is not metric like the Hausdorff distance, but shares most of the properties of a metric. It has been used in many problem on fixed point theory like those in [1, 2, 19, 33].

In recent times, fixed point theory has developed rapidly in partially ordered metric spaces; that is, metric spaces endowed with a partial ordering. References [10, 15, 23, 25, 27] are some recent instances of these works. A speciality of these problems is that they use both analytic and order theoretic methods. It is also one of the main reasons why they are considered interesting.

Khan et al. [21] initiated the use of a control function in metric fixed point theory which they called Altering distance function. Several works on fixed point theory like those noted in [12, 16, 26, 28] have utilized this control function.

---

\*corresponding author

The concept of almost contractions were introduced by Berinde [5, 6]. It was shown in [5] that any strict contraction, the Kannan [20] and Zamfirescu [34] mappings, as well as a large class of quasi-contractions, are all almost contractions. Almost contractions and its generalizations were further considered in several works like [7, 11, 24].

The purpose of this paper is to establish some coincidence point results of a family of multivalued mappings with a single valued mapping under the assumptions of certain almost contractive type inequalities in partially ordered metric spaces. We have also utilized  $\delta$ -compatible pairs in our theorems. In another theorem we have replaced the continuities of the functions with an order condition. We also give here the corresponding singlevalued versions of the theorems which generalize a number of existing works. An illustrative example for the multivalued case is given.

## 2 Mathematical Preliminaries

In the following we give some technical definitions which are used in our theorems. Let  $(X, d)$  be a metric space. We denote the class of nonempty and bounded subsets of  $X$  by  $B(X)$ . For  $A, B \in B(X)$ , functions  $D(A, B)$  and  $\delta(A, B)$  are defined as follows:

$$D(A, B) = \inf \{d(a, b) : a \in A, b \in B\}$$

and

$$\delta(A, B) = \sup \{d(a, b) : a \in A, b \in B\}.$$

If  $A = \{a\}$ , then we write  $D(A, B) = D(a, B)$  and  $\delta(A, B) = \delta(a, B)$ . Also, in addition, if  $B = \{b\}$ , then  $D(A, B) = d(a, b)$  and  $\delta(A, B) = d(a, b)$ . Obviously,  $D(A, B) \leq \delta(A, B)$ . For all  $A, B, C \in B(X)$ , the definition of  $\delta(A, B)$  yields the following:

$$\delta(A, B) = \delta(B, A),$$

$$\delta(A, B) \leq \delta(A, C) + \delta(C, B),$$

$$\delta(A, B) = 0 \text{ iff } A = B = \{a\},$$

$$\delta(A, A) = \text{diam } A. \text{ [13]}$$

There are several works which have utilized  $\delta$  - distance [2, 4, 13, 14, 19, 33].

**Definition 1.** ([13]) A sequence  $\{A_n\}$  of subsets of metric space  $(X, d)$  is said to be convergent to subset  $A$  of  $X$  if

(i) given  $a \in A$ , there is a sequence  $\{a_n\}$  in  $X$  such that  $a_n \in A_n$ , for  $n = 1, 2, 3, \dots$ , and  $\{a_n\}$  converges to  $a$ .

(ii) given  $\epsilon > 0$ , there exists a positive integer  $N$  such that  $A_n \subseteq A_\epsilon$ , for all  $n > N$ , where  $A_\epsilon$  is the union of all open sphere with centers in  $A$  and radius  $\epsilon$ .

**Lemma 2.** ([13, 14]) If  $\{A_n\}$  and  $\{B_n\}$  are sequences in  $B(X)$ , where  $(X, d)$  is a complete metric space and  $\{A_n\} \rightarrow A$  and  $\{B_n\} \rightarrow B$  where  $A, B \in B(X)$  then

$$\delta(A_n, B_n) \rightarrow \delta(A, B) \text{ as } n \rightarrow \infty.$$

**Lemma 3.** ([14]) If  $\{A_n\}$  is a sequence of bounded subsets of a complete metric space  $(X, d)$  and if  $\lim_{n \rightarrow \infty} \delta(A_n, \{y\}) = 0$ , for some  $y \in X$ , then  $\{A_n\} \rightarrow \{y\}$  as  $n \rightarrow \infty$ .

**Definition 4.** ([14]) A set-valued mapping  $T : X \rightarrow B(X)$ , where  $(X, d)$  is a metric space, is continuous at a point  $x \in X$  if  $\{x_n\}$  is a sequence in  $X$  converging to  $x$ , then the sequence  $\{Tx_n\}$  in  $B(X)$  converges to  $Tx$ .  $T$  is said to be continuous in  $X$  if it is continuous at each point  $x \in X$ .

**Lemma 5.** ([14]) If  $\{A_n\}$  is a sequence of nonempty subsets of  $X$  and  $z \in X$  such that

$$\lim_{n \rightarrow \infty} a_n = z,$$

where  $z$  is independent of the particular choice of each  $a_n \in A_n$ . If a self map  $g$  of  $X$  is continuous,  $\{gz\}$  is the limit of the sequence  $\{gA_n\}$ .

**Definition 6.** ([18]) Two self maps  $g$  and  $T$  of a metric space  $(X, d)$  are said to be compatible mappings if  $\lim_{n \rightarrow \infty} d(gTx_n, Tgx_n) = 0$  whenever  $\{x_n\}$  is a sequence in  $X$  such that  $\lim_{n \rightarrow \infty} gx_n = \lim_{n \rightarrow \infty} Tx_n = t$ , for some  $t \in X$ .

**Definition 7.** ([19]) The mappings  $g : X \rightarrow X$  and  $T : X \rightarrow B(X)$ , where  $(X, d)$  is a metric space, are  $\delta$ -compatible if  $\lim_{n \rightarrow \infty} \delta(Tgx_n, gTx_n) = 0$  whenever  $\{x_n\}$  is a sequence in  $X$  such that  $gTx_n \in B(X)$  and  $Tx_n \rightarrow \{t\}$ ,  $gx_n \rightarrow t$ , for some  $t$  in  $X$ .

**Definition 8.** Let  $(X, d)$  be a metric space and  $g : X \rightarrow X$  and  $T : X \rightarrow B(X)$ . Then  $u \in X$  is called a coincidence point of  $g$  and  $T$  if  $\{gu\} = Tu$ .

**Definition 9.** ([4]) Let  $A$  and  $B$  be two nonempty subsets of a partially ordered set  $(X, \preceq)$ . The relation between  $A$  and  $B$  is denoted and defined as follows:

$A \prec_1 B$ , if for every  $a \in A$  there exists  $b \in B$  such that  $a \preceq b$ .

**Definition 10.** ([21]) A function  $\psi : [0, \infty) \rightarrow [0, \infty)$  is called an altering distance function if the following properties are satisfied:

- (i)  $\psi$  is monotone increasing and continuous,
- (ii)  $\psi(t) = 0$  if and only if  $t = 0$ .

### 3 Main Results

**Lemma 11.** Let  $(X, d)$  be a metric space and let  $\{x_n\}$  be a sequence in  $X$  such that

$$\lim_{n \rightarrow \infty} d(x_{n+1}, x_n) = 0. \quad (3.1)$$

If  $\{x_n\}$  is not a Cauchy sequence in  $(X, d)$ , then there exists  $\epsilon > 0$  and two sequences  $\{m(k)\}$  and  $\{n(k)\}$  of positive integers such that  $n(k) > m(k) > k$  and the following four sequences tend to  $\epsilon$  when  $k \rightarrow \infty$ :

$$d(x_{m(k)}, x_{n(k)}), d(x_{m(k)}, x_{n(k)+1}), d(x_{n(k)}, x_{m(k)+1}), d(x_{m(k)+1}, x_{n(k)+1}). \quad (3.2)$$

*Proof.* Suppose that  $\{x_n\}$  is a sequence in  $(X, d)$  satisfying (3.1) which is not Cauchy. Then there exists  $\epsilon > 0$  and two sequences  $\{m(k)\}$  and  $\{n(k)\}$  of positive integers such that for all positive integers  $k$ ,

$$n(k) > m(k) > k, d(x_{m(k)}, x_{n(k)-1}) < \epsilon, d(x_{m(k)}, x_{n(k)}) \geq \epsilon.$$

Now,

$$\epsilon \leq d(x_{m(k)}, x_{n(k)}) \leq d(x_{n(k)}, x_{n(k)-1}) + d(x_{n(k)-1}, x_{m(k)}) < d(x_{n(k)}, x_{n(k)-1}) + \epsilon.$$

Letting  $k \rightarrow \infty$  in the above inequality and using (3.1), we have

$$\lim_{k \rightarrow \infty} d(x_{m(k)}, x_{n(k)}) = \epsilon. \quad (3.3)$$

Again,

$$d(x_{m(k)}, x_{n(k)}) \leq d(x_{m(k)}, x_{n(k)+1}) + d(x_{n(k)+1}, x_{n(k)})$$

and

$$d(x_{m(k)}, x_{n(k)+1}) \leq d(x_{m(k)}, x_{n(k)}) + d(x_{n(k)}, x_{n(k)+1}).$$

Letting  $k \rightarrow \infty$  in the above inequalities and using (3.1) and (3.3), we have

$$\lim_{k \rightarrow \infty} d(x_{m(k)}, x_{n(k)+1}) = \epsilon. \quad (3.4)$$

That the remaining two sequences in (3.2) tend to  $\epsilon$  can be proved in a similar way.  $\square$

**Theorem 12.** Let  $\theta : [0, \infty) \rightarrow [0, 1)$  be a continuous function and  $\psi$  be an altering distance function. Let  $(X, \preceq)$  be a partially ordered set and suppose that there exists a metric  $d$  on  $X$  such that  $(X, d)$  is a complete metric space. Let  $\{T_\alpha : X \rightarrow B(X) : \alpha \in \Lambda\}$  be a family of multivalued mappings. Let  $g : X \rightarrow X$  be a mapping such that  $g(X)$  is closed in  $X$ . Suppose that there exists  $\alpha_0 \in \Lambda$  such that

(i)  $T_{\alpha_0}$  and  $g$  are continuous,

(ii)  $T_{\alpha_0}x \subseteq g(X)$  and  $gT_{\alpha_0}x \in B(X)$ , for every  $x \in X$ ,

(iii) there exists  $x_0 \in X$  such that  $\{gx_0\} \prec_1 T_{\alpha_0}x_0$ ,

(iv) for  $x, y \in X$ ,  $gx \preceq gy$  implies  $T_{\alpha_0}x \prec_1 T_{\alpha_0}y$ ,

(v) the pair  $(g, T_{\alpha_0})$  is  $\delta$ -compatible,

(vi)  $\psi(\delta(T_{\alpha_0}x, T_{\alpha_0}y))$

$$\leq \theta(d(gx, gy)) \max \{ \psi(d(gx, gy)), \psi(D(gx, T_{\alpha_0}x)), \psi(D(gy, T_{\alpha_0}y)),$$

$$\sqrt{\psi(D(gx, T_{\alpha_0}y)) \cdot \psi(D(gy, T_{\alpha_0}x))} \}$$

$$+ L \min \{ \psi(D(gx, T_{\alpha_0}x)), \psi(D(gy, T_{\alpha_0}y)), \psi(D(gx, T_{\alpha_0}y)), \psi(D(gy, T_{\alpha_0}x)) \},$$

where  $x, y \in X$  such that  $gx$  and  $gy$  are comparable and  $L \geq 0$ .

Then  $g$  and  $\{T_\alpha : \alpha \in \Lambda\}$  have a coincidence point.

*Proof.* First we establish that any coincidence point of  $g$  and  $T_{\alpha_0}$  is a coincidence point of  $g$  and  $\{T_\alpha : \alpha \in \Lambda\}$  and conversely. Suppose that  $z \in X$  be a coincidence point of  $g$

and  $T_{\alpha_0}$ . Then  $\{gz\} = T_{\alpha_0}z$ . From (vi) and using the properties of  $\psi$ , we have

$$\begin{aligned} \psi(\delta(gz, T_{\alpha}z)) &= \psi(\delta(T_{\alpha_0}z, T_{\alpha}z)) \\ &\leq \theta(d(gz, gz)) \max \{ \psi(d(gz, gz)), \psi(D(gz, T_{\alpha_0}z)), \psi(D(gz, T_{\alpha}z)), \\ &\quad \sqrt{\psi(D(gz, T_{\alpha}z)) \cdot \psi(D(gz, T_{\alpha_0}z))} \} \\ &+ L \min \{ \psi(D(gz, T_{\alpha_0}z)), \psi(D(gz, T_{\alpha}z)), \psi(D(gz, T_{\alpha}z)), \psi(D(gz, T_{\alpha_0}z)) \} \\ &= \theta(d(gz, gz)) \max \{ \psi(d(gz, gz)), \psi(d(gz, gz)), \psi(D(gz, T_{\alpha}z)), \\ &\quad \sqrt{\psi(D(gz, T_{\alpha}z)) \cdot \psi(d(gz, gz))} \} \\ &+ L \min \{ \psi(d(gz, gz)), \psi(D(gz, T_{\alpha}z)), \psi(D(gz, T_{\alpha}z)), \psi(d(gz, gz)) \} \\ &= \theta(d(gz, gz))\psi(D(gz, T_{\alpha}z)) \\ &< \psi(D(gz, T_{\alpha}z)), \text{ (since } \theta(t) < 1, \text{ for all } t \in [0, \infty)\text{).} \end{aligned}$$

Again using the monotone property of  $\psi$ , we have

$$\delta(gz, T_{\alpha}z) < D(gz, T_{\alpha}z) \leq \delta(gz, T_{\alpha}z),$$

which implies that  $\delta(gz, T_{\alpha}z) = 0$ , that is,  $\{gz\} = T_{\alpha}z$ , for all  $\alpha \in \Lambda$ . Hence  $z$  is a coincidence point of  $g$  and  $\{T_{\alpha} : \alpha \in \Lambda\}$ . Converse part is trivial.

Now it is sufficient to prove that  $g$  and  $T_{\alpha_0}$  have a coincidence point. Let  $x_0 \in X$  be such that  $\{gx_0\} \prec_1 T_{\alpha_0}x_0$ . Then there exists  $u \in T_{\alpha_0}x_0$  such that  $gx_0 \preceq u$ . Since  $T_{\alpha_0}x_0 \subseteq g(X)$  and  $u \in T_{\alpha_0}x_0$ , there exists  $x_1 \in X$  such that  $gx_1 = u$ . So  $gx_0 \preceq gx_1$ . Then by the assumption (iv),  $T_{\alpha_0}x_0 \prec_1 T_{\alpha_0}x_1$ . Since  $u = gx_1 \in T_{\alpha_0}x_0$ , there exists  $v \in T_{\alpha_0}x_1$  such that  $gx_1 \preceq v$ . As  $T_{\alpha_0}x_1 \subseteq g(X)$  and  $v \in T_{\alpha_0}x_1$ , there exists  $x_2 \in X$  such that  $gx_2 = v$ . So  $gx_1 \preceq gx_2$ . Continuing this process we construct a sequence  $\{x_n\}$  in  $X$  such that

$$gx_{n+1} \in T_{\alpha_0}x_n, \text{ for all } n \geq 0, \tag{3.5}$$

and

$$gx_0 \preceq gx_1 \preceq gx_2 \preceq \dots \preceq gx_n \preceq gx_{n+1} \dots \tag{3.6}$$

Let  $\tau_n = d(gx_n, gx_{n+1})$ .

Since  $gx_n \preceq gx_{n+1}$ , putting  $\alpha = \alpha_0$ ,  $x = x_n$  and  $y = x_{n+1}$  in (vi) and using the properties of  $\psi$ , we have

$$\begin{aligned} \psi(\tau_{n+1}) &\leq \psi(\delta(T_{\alpha_0}x_n, T_{\alpha_0}x_{n+1})) \\ &\leq \theta(\tau_n) \max \{ \psi(\tau_n), \psi(D(gx_n, T_{\alpha_0}x_n)), \psi(D(gx_{n+1}, T_{\alpha_0}x_{n+1})), \\ &\quad \sqrt{\psi(D(gx_n, T_{\alpha_0}x_{n+1})) \cdot \psi(D(gx_{n+1}, T_{\alpha_0}x_n))} \} \\ &+ L \min \{ \psi(D(gx_n, T_{\alpha_0}x_n)), \psi(D(gx_{n+1}, T_{\alpha_0}x_{n+1})), \\ &\quad \psi(D(gx_n, T_{\alpha_0}x_{n+1})), \psi(D(gx_{n+1}, T_{\alpha_0}x_n)) \} \\ &\leq \theta(\tau_n) \max \{ \psi(\tau_n), \psi(d(gx_n, gx_{n+1})), \psi(d(gx_{n+1}, gx_{n+2})), \\ &\quad \sqrt{\psi(d(gx_n, gx_{n+2})) \cdot \psi(d(gx_{n+1}, gx_{n+1}))} \} \\ &+ L \min \{ \psi(d(gx_n, gx_{n+1})), \psi(d(gx_{n+1}, gx_{n+2})), \\ &\quad \psi(d(gx_n, gx_{n+2})), \psi(d(gx_{n+1}, gx_{n+1})) \} \\ &= \theta(\tau_n) \max \{ \psi(\tau_n), \psi(\tau_{n+1}) \}. \tag{3.7} \end{aligned}$$

Suppose that,  $\max \{ \psi(\tau_n), \psi(\tau_{n+1}) \} = \psi(\tau_{n+1})$ . Then from (3.7), it follows that

$$\psi(\tau_{n+1}) \leq \theta(\tau_n) \psi(\tau_{n+1}) < \psi(\tau_{n+1}), \text{ (since } \theta(\tau_n) < 1\text{),}$$

which is a contradiction. Hence

$$\psi(\tau_{n+1}) \leq \theta(\tau_n) \psi(\tau_n) < \psi(\tau_n), \quad (\text{since } \theta(\tau_n) < 1).$$

By the monotone property of  $\psi$ , it follows that

$$\tau_{n+1} < \tau_n, \quad \text{for all } n \geq 0,$$

that is,  $\{\tau_n\}$  is a monotone decreasing sequence of nonnegative real numbers. Hence there exists a  $\tau \geq 0$  such that

$$\tau_n \longrightarrow \tau \quad \text{as } n \longrightarrow \infty.$$

Taking  $n \longrightarrow \infty$  in (3.7), using the continuities of  $\theta$  and  $\psi$ , we have

$$\psi(\tau) \leq \theta(\tau) \psi(\tau) < \psi(\tau), \quad (\text{since } \theta(\tau) < 1),$$

which is a contradiction unless  $\tau = 0$ . Thus we have

$$\lim_{n \rightarrow \infty} \tau_n = \lim_{n \rightarrow \infty} d(gx_n, gx_{n+1}) = 0. \quad (3.8)$$

Next we show that  $\{gx_n\}$  is a Cauchy sequence. If  $\{gx_n\}$  is not a Cauchy sequence, then following Lemma 11, there exists  $\epsilon > 0$  and two sequences  $\{m(k)\}$  and  $\{n(k)\}$  of positive integers such that for all positive integers  $k$ ,  $n(k) > m(k) > k$  and

$$\lim_{k \rightarrow \infty} d(gx_{m(k)}, gx_{n(k)}) = \epsilon, \quad (3.9)$$

$$\lim_{k \rightarrow \infty} d(gx_{m(k)}, gx_{n(k)+1}) = \epsilon, \quad (3.10)$$

$$\lim_{k \rightarrow \infty} d(gx_{n(k)}, gx_{m(k)+1}) = \epsilon, \quad (3.11)$$

and

$$\lim_{k \rightarrow \infty} d(gx_{m(k)+1}, gx_{n(k)+1}) = \epsilon. \quad (3.12)$$

For each positive integer  $k$ ,  $gx_{m(k)}$  and  $gx_{n(k)}$  are comparable. Then putting  $\alpha = \alpha_0$ ,  $x = x_{m(k)}$  and  $y = x_{n(k)}$  in (vi) and using the monotone property of  $\psi$ , we have

$$\begin{aligned} \psi(d(gx_{m(k)+1}, gx_{n(k)+1})) &\leq \psi(\delta(T_{\alpha_0}x_{m(k)}, T_{\alpha_0}x_{n(k)})) \\ &\leq \theta(d(gx_{m(k)}, gx_{n(k)})) \max \{ \psi(d(gx_{m(k)}, gx_{n(k)})), \psi(D(gx_{m(k)}, T_{\alpha_0}x_{m(k)})), \\ &\quad \psi(D(gx_{n(k)}, T_{\alpha_0}x_{n(k)})), \\ &\quad \sqrt{\psi(D(gx_{m(k)}, T_{\alpha_0}x_{n(k)})) \cdot \psi(D(gx_{n(k)}, T_{\alpha_0}x_{m(k)}))} \} \\ &\quad + L \min \{ \psi(D(gx_{m(k)}, T_{\alpha_0}x_{m(k)})), \psi(D(gx_{n(k)}, T_{\alpha_0}x_{n(k)})), \\ &\quad \psi(D(gx_{m(k)}, T_{\alpha_0}x_{n(k)})), \psi(D(gx_{n(k)}, T_{\alpha_0}x_{m(k)})) \} \\ &\leq \theta(d(gx_{m(k)}, gx_{n(k)})) \max \{ \psi(d(gx_{m(k)}, gx_{n(k)})), \psi(d(gx_{m(k)}, gx_{m(k)+1})), \\ &\quad \psi(d(gx_{n(k)}, gx_{n(k)+1})), \\ &\quad \sqrt{\psi(d(gx_{m(k)}, gx_{n(k)+1})) \cdot \psi(d(gx_{n(k)}, gx_{m(k)+1}))} \} \\ &\quad + L \min \{ \psi(d(gx_{m(k)}, gx_{m(k)+1})), \psi(d(gx_{n(k)}, gx_{n(k)+1})), \\ &\quad \psi(d(gx_{m(k)}, gx_{n(k)+1})), \psi(d(gx_{n(k)}, gx_{m(k)+1})) \}. \end{aligned}$$

Letting  $k \rightarrow \infty$  in the above inequality, using (3.8), (3.9), (3.10), (3.11) and (3.12) and using the properties of  $\theta$  and  $\psi$ , we have

$$\psi(\epsilon) \leq \theta(\epsilon) \psi(\epsilon) < \psi(\epsilon), \quad (\text{since } \theta(\epsilon) < 1),$$

which is a contradiction. Hence  $\{gx_n\}$  is a Cauchy sequence in  $g(X)$ . Since  $X$  is complete and  $g(X)$  is closed in  $X$ , there exists  $u \in g(X)$  such that

$$gx_n \rightarrow u \quad \text{as } n \rightarrow \infty.$$

Since  $u \in g(X)$ , there exists  $z \in X$  such that  $u = gz$ . Then

$$gx_n \rightarrow u = gz \quad \text{as } n \rightarrow \infty. \quad (3.13)$$

Since  $\{\tau_n\}$  is monotone decreasing, from (3.7), we have

$$\psi(\tau_{n+1}) \leq \psi(\delta(T_{\alpha_0}x_n, T_{\alpha_0}x_{n+1})) \leq \theta(\tau_n)\psi(\tau_n).$$

As  $\theta(\tau_n) < 1$ , it follows that

$$\psi(\tau_{n+1}) \leq \psi(\delta(T_{\alpha_0}x_n, T_{\alpha_0}x_{n+1})) < \psi(\tau_n),$$

which, by the monotone property of  $\psi$ , implies that

$$\tau_{n+1} \leq \delta(T_{\alpha_0}x_n, T_{\alpha_0}x_{n+1}) < \tau_n.$$

Taking  $n \rightarrow \infty$  in the above inequality, and using (3.8), we have

$$\lim_{n \rightarrow \infty} \delta(T_{\alpha_0}x_{n+1}, T_{\alpha_0}x_n) = 0. \quad (3.14)$$

Now,

$$\delta(T_{\alpha_0}x_n, \{u\}) \leq \delta(T_{\alpha_0}x_n, gx_n) + \delta(gx_n, \{u\}) \leq \delta(T_{\alpha_0}x_n, T_{\alpha_0}x_{n-1}) + d(gx_n, u).$$

Taking  $n \rightarrow \infty$  in the above inequality, and using (3.13) and (3.14), we have

$$\lim_{n \rightarrow \infty} \delta(T_{\alpha_0}x_n, \{u\}) = 0,$$

which, by Lemma 3, implies that

$$T_{\alpha_0}x_n \rightarrow \{u\} \quad \text{as } n \rightarrow \infty. \quad (3.15)$$

Since the pair  $(g, T_{\alpha_0})$  is  $\delta$ -compatible, from (3.13) and (3.15), we have

$$\lim_{n \rightarrow \infty} \delta(T_{\alpha_0}gx_n, gT_{\alpha_0}x_n) = 0.$$

As  $g$  and  $T_{\alpha_0}$  are continuous, it follows by Lemma 5 that  $\delta(T_{\alpha_0}u, gu) = 0$ , that is,  $T_{\alpha_0}u = \{gu\}$ . Hence  $u \in g(X) \subseteq X$  is a coincidence point of  $g$  and  $T_{\alpha_0}$ . By what we have already proved,  $u$  is a coincidence point of  $g$  and  $\{T_\alpha : \alpha \in \Lambda\}$ .  $\square$

In our next theorem we relax the continuity assumption on  $T_{\alpha_0}$  and  $g$  by imposing an order condition. We also relax the  $\delta$ -compatibility assumption of the pairs  $(g, T_{\alpha_0})$  and the condition that  $gT_{\alpha_0}x \in B(X)$ , for every  $x \in X$ .

**Theorem 13.** Let  $\theta : [0, \infty) \rightarrow [0, 1)$  be a continuous function and  $\psi$  be an altering distance function. Let  $(X, \preceq)$  be a partially ordered set and suppose that there exists a metric  $d$  on  $X$  such that  $(X, d)$  is a complete metric space. Assume that if  $x_n \rightarrow x$  is a nondecreasing sequence in  $X$ , then  $x_n \preceq x$ , for all  $n$ . Let  $\{T_\alpha : X \rightarrow B(X) : \alpha \in \Lambda\}$  be a family of multivalued mappings. Let  $g : X \rightarrow X$  be a mapping such that  $g(X)$  is closed in  $X$ . Suppose that there exists  $\alpha_0 \in \Lambda$  such that

$$(i) T_{\alpha_0}x \subseteq g(X), \text{ for every } x \in X,$$

$$(ii) \text{ there exists } x_0 \in X \text{ such that } \{gx_0\} \prec_1 T_{\alpha_0}x_0,$$

$$(iii) \text{ for } x, y \in X, gx \preceq gy \text{ implies } T_{\alpha_0}x \prec_1 T_{\alpha_0}y,$$

$$(iv) \psi(\delta(T_{\alpha_0}x, T_{\alpha_0}y))$$

$$\leq \theta(d(gx, gy)) \max \{\psi(d(gx, gy)), \psi(D(gx, T_{\alpha_0}x)), \psi(D(gy, T_{\alpha_0}y)),$$

$$\sqrt{\psi(D(gx, T_{\alpha_0}y)) \cdot \psi(D(gy, T_{\alpha_0}x))}\}$$

$$+ L \min \{\psi(D(gx, T_{\alpha_0}x)), \psi(D(gy, T_{\alpha_0}y)), \psi(D(gx, T_{\alpha_0}y)), \psi(D(gy, T_{\alpha_0}x))\},$$

where  $x, y \in X$  such that  $gx$  and  $gy$  are comparable and  $L \geq 0$ .

Then  $g$  and  $\{T_\alpha : \alpha \in \Lambda\}$  have a coincidence point.

*Proof.* We take the same sequence  $\{gx_n\}$  as in the proof of Theorem 12. Then we have  $gx_{n+1} \in T_{\alpha_0}x_n$ , for all  $n \geq 0$ ,  $\{gx_n\}$  is monotonic nondecreasing and  $gx_n \rightarrow gz$  as  $n \rightarrow \infty$ . Then by the order condition of the metric space, we have  $gx_n \preceq gz$ , for all  $n$ . Using the monotone property of  $\psi$  and the condition (iv), we have

$$\psi(\delta(gx_{n+1}, T_\alpha z)) \leq \psi(\delta(T_{\alpha_0}x_n, T_\alpha z))$$

$$\leq \theta(d(gx_n, gz)) \max \{\psi(d(gx_n, gz)), \psi(D(gx_n, T_{\alpha_0}x_n)), \psi(D(gz, T_\alpha z)),$$

$$\sqrt{\psi(D(gx_n, T_\alpha z)) \cdot \psi(D(gz, T_{\alpha_0}x_n))}\}$$

$$+ L \min \{\psi(D(gx_n, T_{\alpha_0}x_n)), \psi(D(gz, T_\alpha z)), \psi(D(gx_n, T_\alpha z)), \psi(D(gz, T_{\alpha_0}x_n))\}$$

$$\leq \theta(d(gx_n, gz)) \max \{\psi(d(gx_n, gz)), \psi(d(gx_n, gx_{n+1})), \psi(D(gz, T_\alpha z)),$$

$$\sqrt{\psi(D(gx_n, T_\alpha z)) \cdot \psi(d(gz, gx_{n+1}))}\}$$

$$+ L \min \{\psi(d(gx_n, gx_{n+1})), \psi(D(gz, T_\alpha z)), \psi(D(gx_n, T_\alpha z)), \psi(d(gz, gx_{n+1}))\}.$$

Letting  $n \rightarrow \infty$  in the above inequality and using the properties of  $\theta$  and  $\psi$ , we have

$$\psi(\delta(gz, T_\alpha z)) \leq \theta(0)\psi(D(gz, T_\alpha z)) \leq \theta(0)\psi(\delta(gz, T_\alpha z)) < \psi(\delta(gz, T_\alpha z)) \text{ (since } \theta(0) < 1),$$

which implies that  $\delta(gz, T_\alpha z) = 0$ , that is,  $\{gz\} = T_\alpha z$ , for all  $\alpha \in \Lambda$ . Hence  $z$  is a coincidence point of  $g$  and  $\{T_\alpha : \alpha \in \Lambda\}$ .  $\square$

Considering  $\{T_\alpha : X \rightarrow B(X) : \alpha \in \Lambda\} = \{T\}$  in Theorem 12, we have the following corollary.

**Corollary 14.** Let  $\theta : [0, \infty) \rightarrow [0, 1)$  be a continuous function and  $\psi$  be an altering distance function. Let  $(X, \preceq)$  be a partially ordered set and suppose that there exists a

metric  $d$  on  $X$  such that  $(X, d)$  is a complete metric space. Let  $T : X \rightarrow B(X)$  be a multivalued mapping and  $g : X \rightarrow X$  a mapping such that

- (i)  $T$  and  $g$  are continuous,
- (ii)  $Tx \subseteq g(X)$  and  $gTx \in B(X)$ , for every  $x \in X$ , and  $g(X)$  is closed in  $X$ ,
- (iii) there exists  $x_0 \in X$  such that  $\{gx_0\} \prec_1 Tx_0$ ,
- (iv) for  $x, y \in X$ ,  $gx \preceq gy$  implies  $Tx \prec_1 Ty$ ,
- (v) the pair  $(g, T)$  is  $\delta$  - compatible,
- (vi)  $\psi(\delta(Tx, Ty))$

$$\begin{aligned} &\leq \theta(d(gx, gy)) \max \{ \psi(d(gx, gy)), \psi(D(gx, Tx)), \psi(D(gy, Ty)), \\ &\quad \sqrt{\psi(D(gx, Ty)) \cdot \psi(D(gy, Tx))} \} \\ &\quad + L \min \{ \psi(D(gx, Tx)), \psi(D(gy, Ty)), \psi(D(gx, Ty)), \psi(D(gy, Tx)) \}, \end{aligned}$$

where  $x, y \in X$  such that  $gx$  and  $gy$  are comparable and  $L \geq 0$ .

Then  $g$  and  $T$  have a coincidence point.

Considering  $\{T_\alpha : X \rightarrow B(X) : \alpha \in \Lambda\} = \{T\}$  in Theorem 13, we have the following corollary.

**Corollary 15.** Let  $\theta : [0, \infty) \rightarrow [0, 1)$  be a continuous function and  $\psi$  be an altering distance function. Let  $(X, \preceq)$  be a partially ordered set and suppose that there exists a metric  $d$  on  $X$  such that  $(X, d)$  is a complete metric space. Assume that if  $x_n \rightarrow x$  is a nondecreasing sequence in  $X$ , then  $x_n \preceq x$ , for all  $n$ . Let  $T : X \rightarrow B(X)$  be a multivalued mapping and  $g : X \rightarrow X$  a mapping such that

- (i)  $Tx \subseteq g(X)$ , for every  $x \in X$ , and  $g(X)$  is closed in  $X$ ,
- (ii) there exists  $x_0 \in X$  such that  $\{gx_0\} \prec_1 Tx_0$ ,
- (iii) for  $x, y \in X$ ,  $gx \preceq gy$  implies  $Tx \prec_1 Ty$ ,
- (iv)  $\psi(\delta(Tx, Ty))$

$$\begin{aligned} &\leq \theta(d(gx, gy)) \max \{ \psi(d(gx, gy)), \psi(D(gx, Tx)), \psi(D(gy, Ty)), \\ &\quad \sqrt{\psi(D(gx, Ty)) \cdot \psi(D(gy, Tx))} \} \\ &\quad + L \min \{ \psi(D(gx, Tx)), \psi(D(gy, Ty)), \psi(D(gx, Ty)), \psi(D(gy, Tx)) \}, \end{aligned}$$

where  $x, y \in X$  such that  $gx$  and  $gy$  are comparable and  $L \geq 0$ .

Then  $g$  and  $T$  have a coincidence point.

The following theorems are single valued cases of the Theorems 12 and 13 respectively. Here we treat  $T$  as a multivalued mapping in which case  $Tx$  is a singleton set for every  $x \in X$ .

**Theorem 16.** *Let  $\theta : [0, \infty) \rightarrow [0, 1)$  be a continuous function and  $\psi$  be an altering distance function. Let  $(X, \preceq)$  be a partially ordered set and suppose that there exists a metric  $d$  on  $X$  such that  $(X, d)$  is a complete metric space. Let  $\{T_\alpha : X \rightarrow X : \alpha \in \Lambda\}$  be a family of mappings. Let  $g : X \rightarrow X$  be a mapping such that  $g(X)$  is closed in  $X$ . Suppose that there exists  $\alpha_0 \in \Lambda$  such that*

(i)  $T_{\alpha_0}$  and  $g$  are continuous,

(ii)  $T_{\alpha_0}(X) \subseteq g(X)$ ,

(iii) there exists  $x_0 \in X$  such that  $gx_0 \preceq T_{\alpha_0}x_0$ ,

(iv) for  $x, y \in X$ ,  $gx \preceq gy$  implies  $T_{\alpha_0}x \preceq T_{\alpha_0}y$ ,

(v) the pair  $(g, T_{\alpha_0})$  is compatible,

(vi)  $\psi(d(T_{\alpha_0}x, T_{\alpha_0}y))$

$$\leq \theta(d(gx, gy)) \max \{ \psi(d(gx, gy)), \psi(d(gx, T_{\alpha_0}x)), \psi(d(gy, T_{\alpha_0}y)),$$

$$\sqrt{\psi(d(gx, T_{\alpha_0}y)) \cdot \psi(d(gy, T_{\alpha_0}x))} \}$$

$$+ L \min \{ \psi(d(gx, T_{\alpha_0}x)), \psi(d(gy, T_{\alpha_0}y)), \psi(d(gx, T_{\alpha_0}y)), \psi(d(gy, T_{\alpha_0}x)) \},$$

where  $x, y \in X$  such that  $gx$  and  $gy$  are comparable and  $L \geq 0$ .

Then  $g$  and  $\{T_\alpha : \alpha \in \Lambda\}$  have a coincidence point.

**Theorem 17.** *Let  $\theta : [0, \infty) \rightarrow [0, 1)$  be a continuous function and  $\psi$  be an altering distance function. Let  $(X, \preceq)$  be a partially ordered set and suppose that there exists a metric  $d$  on  $X$  such that  $(X, d)$  is a complete metric space. Assume that if  $x_n \rightarrow x$  is a nondecreasing sequence in  $X$ , then  $x_n \preceq x$ , for all  $n$ . Let  $\{T_\alpha : X \rightarrow X : \alpha \in \Lambda\}$  be a family of mappings. Let  $g : X \rightarrow X$  be a mapping such that  $g(X)$  is closed in  $X$ . Suppose that there exists  $\alpha_0 \in \Lambda$  such that*

(i)  $T_{\alpha_0}(X) \subseteq g(X)$ ,

(ii) there exists  $x_0 \in X$  such that  $gx_0 \preceq T_{\alpha_0}x_0$ ,

(iii) for  $x, y \in X$ ,  $gx \preceq gy$  implies  $T_{\alpha_0}x \preceq T_{\alpha_0}y$ ,

(iv)  $\psi(d(T_{\alpha_0}x, T_{\alpha_0}y))$

$$\leq \theta(d(gx, gy)) \max \{ \psi(d(gx, gy)), \psi(d(gx, T_{\alpha_0}x)), \psi(d(gy, T_{\alpha_0}y)),$$

$$\sqrt{\psi(d(gx, T_{\alpha_0}y)) \cdot \psi(d(gy, T_{\alpha_0}x))} \}$$

$$+ L \min \{ \psi(d(gx, T_{\alpha_0}x)), \psi(d(gy, T_{\alpha}y)), \psi(d(gx, T_{\alpha}y)), \psi(d(gy, T_{\alpha_0}x)) \},$$

where  $x, y \in X$  such that  $gx$  and  $gy$  are comparable and  $L \geq 0$ .

Then  $g$  and  $\{T_{\alpha} : \alpha \in \Lambda\}$  have a coincidence point.

**Corollary 18.** Let  $p, q, r, s$  be four continuous functions from  $[0, \infty)$  into  $[0, 1)$  which satisfy the property  $p(t) + q(t) + r(t) + s(t) < 1$ , for all  $t \in [0, \infty)$  and  $\psi$  be an altering distance function. Let  $(X, \preceq)$  be a partially ordered set and suppose that there exists a metric  $d$  on  $X$  such that  $(X, d)$  is a complete metric space. Let  $T : X \rightarrow X$  and  $g : X \rightarrow X$  be two mappings such that

(i)  $T$  and  $g$  are continuous,

(ii)  $T(X) \subseteq g(X)$  and  $g(X)$  is closed in  $X$ ,

(iii) there exists  $x_0 \in X$  such that  $gx_0 \preceq Tx_0$ ,

(iv) for  $x, y \in X$ ,  $gx \preceq gy$  implies  $Tx \preceq Ty$ ,

(v) the pair  $(g, T)$  is compatible,

(vi)  $\psi(d(Tx, Ty))$

$$\leq p(d(gx, gy))\psi(d(gx, gy)) + q(d(gx, gy))\psi(d(gx, Tx)) + r(d(gx, gy))\psi(d(gy, Ty)) \\ + s(d(gx, gy))\sqrt{\psi(d(gx, Ty)) \cdot \psi(d(gy, Tx))},$$

where  $x, y \in X$  such that  $gx$  and  $gy$  are comparable.

Then  $g$  and  $T$  have a coincidence point.

*Proof.* Starting with the inequality (vi), we have

$$\psi(d(Tx, Ty)) \leq p(d(gx, gy))\psi(d(gx, gy)) + q(d(gx, gy))\psi(d(gx, Tx)) \\ + r(d(gx, gy))\psi(d(gy, Ty)) + s(d(gx, gy))\sqrt{\psi(d(gx, Ty)) \cdot \psi(d(gy, Tx))}, \\ \leq \theta(d(gx, gy)) \max\{\psi(d(gx, gy)), \psi(d(gx, Tx)), \psi(d(gy, Ty)) \\ \sqrt{\psi(d(gx, Ty)) \cdot \psi(d(gy, Tx))}\},$$

where  $\theta(d(gx, gy)) = p(d(gx, gy)) + q(d(gx, gy)) + r(d(gx, gy)) + s(d(gx, gy))$ , which is a special case of the inequality (vi) of Theorem 16 obtained by considering  $\{T_{\alpha} : X \rightarrow X : \alpha \in \Lambda\} = \{T\}$  and  $L = 0$ .  $\square$

**Corollary 19.** Let  $p, q, r, s$  be four continuous functions from  $[0, \infty)$  into  $[0, 1)$  which satisfy the property  $p(t) + q(t) + r(t) + s(t) < 1$ , for all  $t \in [0, \infty)$  and  $\psi$  be an altering distance function. Let  $(X, \preceq)$  be a partially ordered set and suppose that there exists a metric  $d$  on  $X$  such that  $(X, d)$  is a complete metric space. Assume that if  $x_n \rightarrow x$  is a nondecreasing sequence in  $X$ , then  $x_n \preceq x$ , for all  $n$ . Let  $T : X \rightarrow X$  and  $g : X \rightarrow X$  be two mappings such that

(i)  $T(X) \subseteq g(X)$  and  $g(X)$  is closed in  $X$ ,

(ii) there exists  $x_0 \in X$  such that  $gx_0 \preceq Tx_0$ ,

(iii) for  $x, y \in X$ ,  $gx \preceq gy$  implies  $Tx \preceq Ty$ ,

(iv)  $\psi(d(Tx, Ty))$

$$\leq p(d(gx, gy))\psi(d(gx, gy)) + q(d(gx, gy))\psi(d(gx, Tx)) + r(d(gx, gy))\psi(d(gy, Ty)) \\ + s(d(gx, gy))\sqrt{\psi(d(gx, Ty)) \cdot \psi(d(gy, Tx))},$$

where  $x, y \in X$  such that  $gx$  and  $gy$  are comparable.

Then  $g$  and  $T$  have a coincidence point.

*Proof.* Like the proof of the Corollary 18, we can show that the inequality (iv) is a special case of the inequality (iv) of Theorem 17 obtained by considering  $\{T_\alpha : X \rightarrow X : \alpha \in \Lambda\} = \{T\}$  and  $L = 0$ .  $\square$

**Example 20.** Let  $X = [1, \infty)$  with usual order  $\preceq$  be a partially ordered set. Let  $d : X \times X \rightarrow \mathbb{R}$  be given as

$$d(x, y) = |x - y|, \text{ for } x, y \in X.$$

Then  $(X, d)$  is a complete metric space with the required properties mentioned in Theorems 12 and 13.

Let  $g : X \rightarrow X$  be defined as follows:

$$gx = x^2, \text{ for } x \in X.$$

Then  $g$  has the properties mentioned in Theorems 12 and 13.

Let  $\Lambda = \{1, 2, 3, \dots\}$ . Let the family of mappings  $\{T_\alpha : X \rightarrow B(X) : \alpha \in \Lambda\}$  be defined as follows:

$$T_1x = \{1\}, \text{ for } x \in X \quad \text{and for } \alpha \geq 2, \quad T_\alpha x = \begin{cases} \{1\}, & \text{if } 1 \leq x \leq 4, \\ \{1, \frac{2\alpha}{\alpha+1}\}, & \text{if } x > 4. \end{cases}$$

For any sequence  $\{x_n\}$  in  $X$ ,  $T_1x_n \rightarrow \{t\}$ ,  $gx_n \rightarrow t$ , for some  $t$  in  $X$  implies  $t = 1$ . Then clearly, the pair  $(g, T_1)$  is  $\delta$ -compatible. Also,  $g$  and  $T_1$  satisfy required conditions mentioned in Theorems 12 and 13.

Let  $\psi : [0, \infty) \rightarrow [0, \infty)$  be defined as follows:

$$\psi(t) = t^2, \text{ for } t \in [0, \infty).$$

Then  $\psi$  has the properties mentioned in Theorems 12 and 13.

Let  $\theta : [0, \infty) \rightarrow [0, 1)$  be defined as follows:

$$\theta(t) = \frac{1}{2}, \text{ for all } t \in [0, \infty).$$

Then  $\theta$  satisfies the required properties mentioned in Theorems 12 and 13.

The condition (vi) of Theorem 12 and the condition (iv) of Theorem 13 are satisfied for any  $L \geq 0$ . Hence all the condition of Theorems 12 and 13 are satisfied and it is seen that 1 is a coincidence point of  $g$  and  $\{T_\alpha : \alpha \in \Lambda\}$ .

**Note** In the above example if one takes  $g : X \rightarrow X$  to be function as follows:

$$gx = \begin{cases} \frac{x}{2}, & \text{if } 1 \leq x \leq 4, \\ 200, & \text{if } x > 4. \end{cases}$$

Then the above example is still applicable to Theorem 13 but not applicable to Theorem 12 because  $g$  is not continuous and hence does not satisfy required properties mentioned in Theorem 12.

**Remark 21.** Theorems 16 and 17 are generalizations of ordered versions of theorem 3.1 in [8] which generalizes the Banach contraction principle [3], theorem 2 of Khan et al [21], the theorem of Skof [32], and the theorem of Kannan [20]. Also, Theorems 16 and 17 generalize the ordered versions of the main result of Berinde [5].

### Acknowledgement

The authors gratefully acknowledge the suggestions made by the learned referee.

### References

- [1] M. A. Ahmed, Common fixed point theorems for weakly compatible mappings, *Rocky Mountain J. Math.* **33** (2003), 1189–1203.
- [2] I. Altun and D. Turkoglu, Some fixed point theorems for weakly compatible multivalued mappings satisfying an implicit relation, *Filomat* **22** (2008), 13–21.
- [3] S. Banach, Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales, *Fund. Math.* **3** (1922), 133–181.
- [4] I. Beg and A. R. Butt, Common fixed point for generalized set valued contractions satisfying an implicit relation in partially ordered metric spaces, *Math. Commun.* **15** (2010), 65–76.
- [5] V. Berinde, Approximating fixed points of weak contractions using the Picard iteration, *Nonlinear Anal. Forum* **9** (2004), 43–53.
- [6] V. Berinde, General constructive fixed point theorems for Ćirić-type almost contractions in metric spaces, *Carpathian J. Math.* **24** (2008), 10–19.
- [7] V. Berinde, Common fixed points of noncommuting almost contractions in cone metric spaces, *Math. Commun.* **15** (2010), 229–241.
- [8] I. Bhaumik, K. Das, N. Metiya and B. S. Choudhury, A Coincidence point result by using altering distance function, *J. Math. Comput. Sci.* **2** (2012), 61–72.
- [9] H. Bouhadjera, A. Djoudi and B. Fisher, A unique common fixed point theorem for occasionally weakly compatible maps, *Surveys Math. Appl.* **3** (2008), 177–182.
- [10] B. S. Choudhury and A. Kundu,  $(\psi, \alpha, \beta)$  - Weak contractions in partially ordered metric spaces, *Appl. Math. Lett.* **25** (2012), 6–10.
- [11] L. Ćirić, M. Abbas, R. Saadati and N. Hussain, Common fixed points of almost generalized contractive mappings in ordered metric spaces, *Appl. Math. Comput.* **217** (2011), 5784–5789.
- [12] P. N. Dutta and B. S. Choudhury, A generalisation of contraction principle in metric spaces, *Fixed Point Theory Appl.* **2008** (2008), Article ID 406368, 8 pages.
- [13] B. Fisher, Common fixed points of mappings and setvalued mappings, *Rostock Math. Colloq.* **18** (1981), 69–77.
- [14] B. Fisher and S. Sessa, Two common fixed point theorems for weakly commuting mappings, *Period. Math. Hungar.* **20** (1989), 207–218.

- [15] Z. Golubović, Z. Kadelburg and S. Radenović, Common fixed points of ordered  $g$ -quasicontractions and weak contractions in ordered metric spaces, *Fixed Point Theory Appl.* 2012, 2012 : 20.
- [16] J. Harjani and K. Sadarangani, Generalized contractions in partially ordered metric spaces and applications to ordinary differential equations, *Nonlinear Anal.* **72** (2010), 1188–1197.
- [17] S. Hong, Fixed points of multivalued operators in ordered metric spaces with applications, *Nonlinear Anal.* **72** (2010), 3929–3942.
- [18] G. Jungck, Compatible mappings and common fixed points, *Inst. J. Math. Sci.* **9** (1986), 771–779.
- [19] G. Jungck and B. E. Rhoades, Some fixed point theorems for compatible maps, *Inter. J. Math. Sci.* **16** (1993), 417–428.
- [20] R. Kannan, Some results on fixed points, *Bull. Calcutta Math. Soc.* **10** (1968), 71–76.
- [21] M. S. Khan, M. Swaleh and S. Sessa, Fixed points theorems by altering distances between the points, *Bull. Austral. Math. Soc.* **30** (1984), 1–9.
- [22] S. B. Nadler Jr., Multivalued contraction mappings, *Pacific J. Math.* **30** (1969), 475–488.
- [23] J. J. Nieto and R. López, Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations, *Order* **22** (2005), 223–239.
- [24] M. Pacurar, Fixed point theory for cyclic Berinde operators, *Fixed Point Theory* **12** (2011), 419–428.
- [25] S. Radenović and Z. Kadelburg, Generalized weak contractions in partially ordered metric spaces, *Comput. Math. Appl.* **60** (2010), 1776–1783.
- [26] S. Radenović, Z. Kadelburg, D. Jandrlić and A. Jandrlić, Some results on weakly contractive maps, *Bull. Iranian Math. Soc.*, 2011 (In Press).
- [27] A. C. M. Ran and M. C. B. Reurings, A fixed point theorem in partially ordered sets and some applications to matrix equations, *Proc. Amer. Math. Soc.* **132** (2004), 1435–1443.
- [28] K. P. R. Sastry and G. V. R. Babu, Some fixed point theorems by altering distances between the points, *Indian J. Pure Appl. Math.* **30** (1999), 641–647.
- [29] M. Shen and S. Hong, Common fixed points for generalized contractive multivalued operators in complete metric spaces, *Appl. Math. Lett.* **22** (2009), 1864–1869.
- [30] S. Shukla and R. Sen, Set-valued Prešić-Reich type mappings in metric spaces, *RACSAM*, 2012, Doi- 10.1007/s13398-012-0114-2.
- [31] S. Shukla, R. Sen and S. Radenović, Set-valued Prešić type contraction in metric spaces, *Annals of the Allexandru Ioan Cuza University-Mathematics*, 2012 (In Press).
- [32] F. Skof, Teorema di punti fisso per applicazioni negli spazi metrici, *Atti. Accad. Sci. Torino.* **111** ( 1977), 323–329.
- [33] J. Yin and T. Guo, Some fixed point results for a class of  $g$ -monotone increasing multivalued mappings, *Arab J. Math. Sci.* **19** (2013), 35–47.
- [34] T. Zamfirescu, Fixed point theorems in metric spaces, *Arch. Mat. (Basel)* **23** (1972), 292–298.